

Branch points of substitutions and closing ordered Bratteli diagrams [†]

Reem Yassawi

Dept. of Mathematics, Trent University
1600 West Bank Drive, Peterborough, Ontario, K9J 7B8, Canada.
 (e-mail: ryassawi@trentu.ca)

Abstract. We study stationary ordered Bratteli diagrams and give necessary and sufficient conditions for these orders to generate a continuous Vershik map. We apply this to finding adic representations for one sided substitution subshifts. We give an algorithm to find the branch points of a substitution, which have to be mapped to the minimal elements of such an ordering. We find adic representations for substitutions with one branch point, and also substitutions all of whose branch points are fixed.

1. Introduction

While adic representations of primitive two-sided substitution subshifts have been studied in [For97] and [DHS99], and generalised to aperiodic two-sided substitution subshifts in [BKM09], this article originated with the question of whether such a representation existed for one-sided substitution subshifts. This question is a special case of the general question of how far two-sided concepts and results in [HPS92] and [GPS95] about minimal homeomorphisms of a Cantor space can be generalised to minimal continuous maps of a Cantor space. Conjugacy of a pair of two-sided subshifts does not imply conjugacy of the corresponding one-sided subshifts (see Example 14 for an example where the subshifts are generated by substitutions).

An example of a markedly different situation in the one-sided case is the generalisation of Theorem 4.6 in [HPS92], which tells us that any minimal homeomorphism (X, T) of a Cantor space has an adic representation. The proof of this result requires a sequence of refining, generating clopen partitions $(\mathcal{P}_n)$, which naturally define an ordered Bratteli diagram $\mathcal{B}$. If the intersection of the bases of $\mathcal{P}_n$ is a unique point, then the ordering is *proper*, so that the Vershik-adic map $V_{\mathcal{B}} : X_{\mathcal{B}} \rightarrow X_{\mathcal{B}}$ representing T is a homeomorphism. For a continuous minimal map $T : X \rightarrow X$ of a Cantor space, first there is the question of finding the non-invertible points for T - we call these *branch points*; second we need a sequence of refining, generating clopen partitions the intersection of whose bases is the branch points. In addition though the ordered Bratteli diagram generated by this sequence of partitions must have the correct number of maximal paths, and must also support a continuous adic map, so, in the case where there are n branch points, there is the question of determining when an ordered Bratteli diagram with n minimal paths and N maximal paths has a continuous Vershik map defined on it - we call such ordered diagrams *closing*.

[†] This research was partially supported by an NSERC grant.

For (aperiodic and primitive) substitution subshifts (X_τ, σ) -see Section 2 for definitions - the natural sequence of generating partitions $(\mathcal{P}_n)$ to consider is that where $\mathcal{P}_n$ is generated by the τ^n -words. The resulting ordered Bratteli diagram $\mathcal{B}_\tau$ has as many minimal elements as there are right fixed τ -fixed points $x_0x_1 \dots$ in $X_\tau^\mathbb{N}$ and as many maximal elements as there are left fixed points $\dots x_{-2}x_{-1}$ in $X_\tau^\mathbb{N}$. The proof in [DHS99] of the statement that every two sided (aperiodic, primitive) substitution subshift $(X_\tau^\mathbb{Z}, \sigma)$ is topologically conjugate to some adic system (X_B, V_B) consists then of finding a substitution τ' such that $(X_\tau^\mathbb{Z}, \sigma) \cong (X_{\tau'}^\mathbb{Z}, \sigma)$ and where τ' is *proper* -ie has exactly one of each of these left and right τ -fixed points.

Suppose $(X_\tau^\mathbb{N}, \sigma)$ has n branch points $\mathbf{x}_1, \dots, \mathbf{x}_n$ where $\mathbf{x}_i$ is a k_i -branch point, (so that $\mathbf{x}_i$ has k_i σ -preimages). Finding an adic representation for $(X_\tau^\mathbb{N}, \sigma)$, the one-sided subshift defined by τ , requires finding τ' such that

$$(X_{\tau'}^\mathbb{N}, \sigma) \stackrel{\Phi}{\cong} (X_\tau^\mathbb{N}, \sigma), \quad (1)$$

$$\tau' \text{ has } n \text{ right fixed points } \{\mathbf{v}_i\}_{i=1}^n \text{ with } \Phi(\mathbf{x}_i) = \mathbf{v}_i \text{ for } 1 \leq i \leq n, \quad (2)$$

$$\tau' \text{ has } \sum_{i=1}^n k_i \text{ left fixed points } \{\mathbf{u}_j^i : 1 \leq j \leq k_i, 1 \leq i \leq n\}, \quad (3)$$

and

$$u_j^i v_{i'} \in \mathcal{L}_{\tau'} \text{ if and only if } i = i'. \quad (4)$$

Here $\mathcal{L}_{\tau'}$ is the language of τ' , and the right (left) fixed point $\mathbf{u}$ is generated by the letter u whose substitution word has u as a prefix (suffix). Condition 2 is a requirement that branch points of τ are mapped to fixed points of τ' , so that the sequence of generating partitions defined by τ' have bases whose intersection is $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$, and also span the topology of $X_\tau^\mathbb{N}$, (Proposition 8). Condition 3 ensures that an adic representation of τ' has the right number of maximal elements. Condition 4 is the key to ensuring that the ordered diagram $\mathcal{B}_{\tau'}$ is closing. This is explained in Proposition 9.

In Section 2 we lay out definitions and state standard results on substitutions that will be used. We assume throughout that τ is aperiodic, recognizable and all of its powers are injective on letters. Apart from Example 12, we assume τ is primitive. These are standard assumptions so that Proposition 8 applies. An algorithm which finds branch points of a substitution is described completely in Section 3. We remark that branch points are the one-sided counterpart to the *left (or right) asymptotic* points in a two-sided subshift, which have been studied, for example, in [HZ01] and [BDH03] - two bi-infinite points are right asymptotic if they agree from some index N onwards - for us the right infinite ray of the N -th shift of these points is a branch point. We characterise non- τ -fixed left branch points $\mathbf{y}$ of τ as points which satisfy an equation of the form $\mathbf{w}\tau(\mathbf{y}) = \mathbf{y}$ for some non-empty finite word $\mathbf{w}$ (Lemma 6). If $\mathbf{u}$ is a τ -fixed point, Proposition 4 characterises when $\mathbf{u}$ is a branch point.

In Section 4 we define stationary Bratteli diagrams and prove Proposition 9, which characterises in terms of $\mathcal{L}_\tau$ when an adic map on a stationary diagram B with an ordering generated by τ is continuous. This result has a generalisation to arbitrary finite rank ordered diagrams and this is the basis for most of the results in [BKMY11].

In Section 5 we deal with the special case where the substitution subshift $(X_\tau^\mathbb{N}, \sigma)$ has a unique branch point, and call these subshifts *quasi-invertible*. Subshifts with this property have been studied: Sturmian sequences, and more generally, Arnoux-Rauzy sequences ([AR91]) generate quasi-invertible subshifts - the condition (*) in [AR91] ensures this, and in many cases these sequences are fixed points of substitutions (see [CMPS93] for

a characterization of when Sturmian subshifts are substitution subshifts; if the ‘S-adic’ expansion of a sequence in [AR91] is periodic, then it is the fixed point of a substitution, for example the tribonacci substitution). All these subshifts have nice geometric representations as toral rotations, or interval exchange maps. Many (but not all) Pisot substitutions that have been studied in the literature are quasi-invertible. Any substitution with a rational Perron eigenvalue has the two-sided subshift $(X_{\tau}^{\mathbb{Z}}, \sigma)$ orbit equivalent to $(X_{\tau'}^{\mathbb{Z}}, \sigma)$ where τ' is quasi-invertible (see Example 13). It would be interesting to study whether quasi-invertibility imposes any spectral constraints. While not all have discrete spectrum (see Example 12: the Chacon substitution is quasi-invertible) spectral multiplicity may be bounded. If τ is quasi-invertible, then Condition (4) is immediately satisfied, and that a substitution satisfying Conditions (1), (2) and (3) exists is the statement of Theorems 13 and 18. We remark that a τ' satisfying Conditions (1) and (2) for constant length substitutions on a two-letter alphabet is found in Section 7 of [CK71].

If τ has several branch points, all of which are fixed, then we prove in Theorem 21 that the right τ' can be found. For general τ we are not able to find the right substitutions that satisfy Condition 2 and we pose the question of whether given a τ , $(X_{\tau}^{\mathbb{N}}, \sigma) \cong (X_{\tau'}^{\mathbb{N}}, \sigma)$ where all of τ ’s branch points are mapped to τ' -fixed points.

The author thanks Ian Putnam and Fabien Durand for helpful discussions.

2. Preliminaries

2.1. Notation Let $\mathcal{A}$ be a finite alphabet. A *word* $\mathbf{w}$ (denoted using boldface) from $\mathcal{A}$ is a finite concatenation of elements from $\mathcal{A}$. Let $\mathcal{A}^+$ denote the set of words (including the empty word) from $\mathcal{A}$. If $\mathbf{w} = w_1 \dots w_m$ and $\mathbf{w}^* = w_1^* \dots w_n^*$, then $\mathbf{w}\mathbf{w}^* := w_1 \dots w_m w_1^* \dots w_n^*$. Define $\mathbf{w}_{[i,j]} = w_i w_{i+1} \dots w_j$. If $\mathbf{w} = w_1 \dots w_m$, the *length* of $\mathbf{w}$, written $|\mathbf{w}|$, is m . Given words $\mathbf{w}, \mathbf{w}^*$, say that $\mathbf{w}$ *occurs* in $\mathbf{w}^*$, or $\mathbf{w}$ *is a subword of* $\mathbf{w}^*$, if $\mathbf{w} = \mathbf{w}^*_{[i, i+|\mathbf{w}|-1]}$ for some i . If $|\mathbf{w}| = m$, define the k -*shift* $\sigma^k(\mathbf{w}) = \mathbf{w}_{[k+1, m]}$. $\mathbf{w}^*$ is a *prefix* of $\mathbf{w}$ if $\mathbf{w}_{[1, |\mathbf{w}^*|]} = \mathbf{w}^*$, similarly $\mathbf{w}^*$ is a *suffix* of $\mathbf{w}$ if $\mathbf{w}_{[|\mathbf{w}|-|\mathbf{w}^*|+1, |\mathbf{w}|]} = \mathbf{w}^*$. When we speak of ‘the’ prefix (suffix) of a word $\mathbf{w}$, we will mean the first (last) letter of $\mathbf{w}$.

Let $\mathbb{N} := \{0, 1, \dots\}$. If $\mathbb{M} = \mathbb{Z}$ or $\mathbb{N}$, then the space of all $\mathbb{M}$ -indexed sequences from $\mathcal{A}$ is written as $\mathcal{A}^{\mathbb{M}}$, and a configuration $\mathbf{x} \in \mathcal{A}^{\mathbb{M}}$ is written $\mathbf{x} = (x_m)_{m \in \mathbb{M}}$. Let $\mathcal{A}$ be endowed with the discrete topology and $\mathcal{A}^{\mathbb{M}}$ with the product topology (which is metrizable); if $b \in \mathcal{A}$ and $j \in \mathbb{M}$, the clopen sets $[b]_j := \{\mathbf{x} : x_j = b\}$ form a countable basis for the topology on $\mathcal{A}^{\mathbb{M}}$ -if $j = 0$; we write $[b]$ instead of $[b]_0$. The (left) *shift map* $\sigma : \mathcal{A}^{\mathbb{M}} \rightarrow \mathcal{A}^{\mathbb{M}}$ is the map defined as $(\sigma(\mathbf{x}))_m = x_{m+1}$. (X, σ) is a *subshift of* $(\mathcal{A}^{\mathbb{M}}, \sigma)$ if X is a closed σ -invariant subset of $\mathcal{A}^{\mathbb{M}}$.

2.2. Substitutions A *substitution* is a map $\tau : \mathcal{A} \rightarrow \mathcal{A}^+$. We extend τ to a map $\tau : \mathcal{A}^+ \rightarrow \mathcal{A}^+$ by concatenation: if $\mathbf{a} = a_1 \dots a_k$, then $\tau(\mathbf{a}) := \tau(a_1) \dots \tau(a_k)$. In this way iteration τ^n is well defined. The substitution τ is extended to a map $\tau : \mathcal{A}^{\mathbb{N}} \rightarrow \mathcal{A}^{\mathbb{N}}$ defined by $\tau(\mathbf{x}) := \tau(x_0) \tau(x_1) \dots$, and also $\tau : \mathcal{A}^{\mathbb{Z}} \rightarrow \mathcal{A}^{\mathbb{Z}}$ defined by $\tau(\mathbf{x}) := \dots \tau(x_{-1}) \cdot \tau(x_0) \tau(x_1) \dots$. We say τ is *left (right) proper* if there exists l in $\mathcal{A}$ such that l is the prefix (suffix) of $\tau(a)$ for each $a \in \mathcal{A}$. If τ is both left and right proper, it is called *proper*. We say τ *suffix-permutative* if the set of suffixes of $\{\tau(a)\}_{a \in \mathcal{A}}$ is a permutation of $\mathcal{A}$. Also τ is *injective* if $\tau(a) \neq \tau(b)$ whenever $a \neq b$. If $|\mathcal{A}| = d$, then let $M = (m_{i,j})_{1 \leq i, j \leq d}$ be the *incidence matrix* for τ , where $m_{i,j}$ is the number of occurrences of j in $\tau(i)$. We say τ is *primitive* if there exists a positive integer k such that for any $a \in \mathcal{A}$, all letters of $\mathcal{A}$ appear in $\tau^k(a)$, ie some power of the incidence matrix is positive. Let $\mathcal{L}_{\tau}$ be the language generated by the words $\{\tau^n(a) : n \in \mathbb{N}^+, a \in \mathcal{A}\}$. A (right infinite) *fixed point* of τ is a sequence $\mathbf{u} \in \mathcal{A}^{\mathbb{N}}$ such

that $\tau(\mathbf{u}) = \mathbf{u}$. If $\mathbf{u} \in \mathcal{A}^{\mathbb{N}}$ is a fixed point for τ , then $\tau(u_0)$ starts with u_0 . Conversely, if there exists a letter l such that l is a prefix of $\tau(l)$, then $\mathbf{u} := \lim_{n \rightarrow \infty} \tau^n(l)$ is the unique fixed point satisfying $u_0 = l$. Similarly one can define a *left infinite* fixed point using a letter u_0 such that $\tau(u_0)$ ends with u_0 . Using the pigeonhole principle, there exists $n \geq 1$ such that τ^n has at least one fixed point. If τ is primitive, then any fixed point $\mathbf{u}$ generates the same subshift $X_{\tau}^{\mathbb{N}} := \overline{\{\sigma^n(\mathbf{u})\}}$. Henceforth we will be working mainly with one sided substitution subshifts, and unless otherwise indicated, will assume this, and write (X_{τ}, σ) instead of $(X_{\tau}^{\mathbb{N}}, \sigma)$.

In what follows we will almost always assume that our substitution τ is aperiodic and primitive, the exception being in Example 12. There τ has a fixed point $\mathbf{u}$ which generates $\mathcal{L}_{\tau}$ (we call this a *generating* fixed point) and the resulting subshift (X_{τ}, σ) is still minimal (ie for each x in X_{τ} , $\{\sigma^n(x) : n \geq 0\}$ is dense in X_{τ}). Such substitutions enjoy many established properties that primitive substitutions do, and minor modifications of proofs of facts for primitive substitutions can be made to ensure that the same results hold for these substitutions -see [BKM09].

Unilateral recognizability was a condition introduced in [Hos86] and [Que87] which ensured that one could uniquely ‘de-substitute’ sequences in X_{τ} , and find clopen generating partitions for (X_{τ}, σ) . Specifically, the set of (1)-cuttings of $\mathbf{u}$ are

$$E := \{0\} \cup \{|\tau(\mathbf{u}_{[0, p-1]})| : p > 0\}.$$

We then say that τ is (*unilaterally*) *recognizable* if there exists some L so that if $\mathbf{u}_{[i, i+L-1]} = \mathbf{u}_{[j, j+L-1]}$ and $i \in E$, then $j \in E$. A word $\mathbf{w}$ occurring at $\mathbf{u}_i$ and $\mathbf{u}_j$ has the *same 1-cutting* at i and j , with $i < j$, if $E \cap \{i, \dots, i + |\mathbf{w}|\} = j - i + E \cap \{j, \dots, j + |\mathbf{w}|\}$.

The necessary and sufficient conditions that exist in Theorem 3.1, [Mos92] for a primitive substitution to be unilaterally recognizable can be modified (see [BKM09]) to work for substitutions:

THEOREM 1. *Let τ be minimal with generating fixed point $\mathbf{u}$. τ is not recognizable if and only if for each L , there exists a word $\mathbf{w}$ of length L , and two elements $a, b \in \mathcal{A}$ such that*

1. $\tau(b)$ is a proper suffix of $\tau(a)$, and
2. The words $\tau(a)\mathbf{w}$ and $\tau(b)\mathbf{w}$ appear in $\mathbf{u}$ with the same 1-cutting of $\mathbf{w}$.

■

In particular, suffix permutative substitutions are recognizable. It is straightforward to show that if τ is recognizable, and injective, then τ^n is recognizable for each n .

In Lemmas 2 and 3, [Hos86], the following is shown for primitive recognizable substitutions (a similar proof works if τ is minimal). Note that $\tau(X_{\tau}) \subset X_{\tau}$.

PROPOSITION 2. *Let τ be a primitive, recognizable substitution with generating fixed point $\mathbf{u}$.*

1. $\mathbf{y} \in \tau(X_{\tau})$ if and only if $\sigma^{n_i} \mathbf{u} \rightarrow \mathbf{y}$ where $n_i \in E$ for all large i .
2. $\tau(X_{\tau})$ is clopen in X_{τ} .
3. $\sigma^p(\tau(x)) \in \tau(X_{\tau})$ if and only if $p = |\tau(\mathbf{x}_{[0, r]})|$ for some r so that $\sigma^p(\tau(x)) = \tau(\sigma^{r+1}(\mathbf{x}))$.

If τ is also injective, then

4. $\tau : X_{\tau} \rightarrow \tau(X_{\tau})$ is a homeomorphism.

■

We remark here that the previous proposition also implies that for each $n \geq 1$, $\tau^n : X_{\tau} \rightarrow \tau^n(X_{\tau})$ is also a homeomorphism, though as a substitution τ^n is not necessarily injective.

Henceforth we assume that our substitution τ is recognizable and all powers of τ are injective, though the latter assumption is not required for the section on branch points.

COROLLARY 3. Every $\mathbf{y} \in \tau(X_\tau)$ can be written in a unique way as $\mathbf{y} = \tau(\mathbf{x})$ with $\mathbf{x} \in X_\tau$. Every $\mathbf{y} \in X_\tau \setminus \tau(X_\tau)$ can be written as $\mathbf{y} = \sigma^k \tau(\mathbf{x})$ where $\sigma(\mathbf{x})$ is unique, and $0 < k < |\tau(x_0)|$.

Proof: The first assertion follows immediately from Proposition 2. If $\mathbf{y} \in X_\tau \setminus \tau(X_\tau)$ then by Part 1 of Proposition 2, there is a sequence $\{n_i\}$ of integers not in E , with $\sigma^{n_i}(\mathbf{u}) \rightarrow \mathbf{y}$. Now each $n_i = k_i + r_i$ where $k_i \in E$, and the positive r_i is strictly less than the successor of k_i in E . Since E is almost periodic, by dropping to a subsequence if necessary, we assume that there is some fixed positive r such that each $n_i = k_i + r$. Now we can assume that $\sigma^{k_i}(\mathbf{u})$ converges to some $\mathbf{x} \in \tau(X_\tau)$. So $\mathbf{y} = \sigma^r \mathbf{x}$.

Now suppose that $\mathbf{y} = \sigma^r \mathbf{x} = \sigma^{r'} \mathbf{x}'$, where if $\mathbf{x} = x_0 x_1 \dots$ then $r < |\tau(x_0)|$, and similarly for r' . Let $l = |\tau(x_0)| - r$, and define l' similarly.

If $l = l'$, then $\sigma^{r+l} \tau(\mathbf{x}) = \tau(\sigma \mathbf{x}) \in \tau(X_\tau)$ and similarly $\sigma^{r'+l'} \tau(\mathbf{x}') = \tau(\sigma \mathbf{x}') \in \tau(X_\tau)$, so $\tau(\sigma \mathbf{x}) = \tau(\sigma \mathbf{x}')$. Thus $\sigma \mathbf{x} = \sigma \mathbf{x}'$.

Finally we will show that $l \neq l'$ leads to a contradiction. Suppose $l < l'$. Then $\sigma^{r+l} \tau(\mathbf{x}) = \tau(\sigma \mathbf{x}) \in \tau(X_\tau)$, and so $\sigma^{r'+l} \tau(\mathbf{x}') = \sigma^{r+l} \tau(\mathbf{x}) \in \tau(X_\tau)$. Thus by part 3 of Proposition 2, $r' + l = |\tau(\mathbf{x}'_{[0,j]})|$ for some j . This contradicts $r' + l < r' + l' = |\tau(x'_0)|$. ■

3. Branch points

If (Y, T) is a system where $\mathbf{y}$ is not T -invertible, we call $\mathbf{y}$ a *branch point*; if $\mathbf{y}$ has M pre-images under T , we say that $\mathbf{y}$ is an M -branch point. Let us call systems (Y, T) with only one branch point *quasi-invertible*. If (X_τ, σ) is quasi-invertible, and the branch point is an M -branch point, we'll call τ *M-quasi-invertible*. More generally, if a substitution τ has n τ -right fixed points $\{\mathbf{x}_1, \dots, \mathbf{x}_n\}$ where $\mathbf{x}_i$ is a k_i -branch point and $N := \sum_{i=1}^n k_i$, call τ an (N, n) -substitution. In this section we describe the substitution structure of branch points $\mathbf{x} \in X_\tau$, and describe an algorithm which finds all such points. Say that a_1 is *part of a suffix cycle* $a_1, a_2, \dots, a_m$ if $\tau(a_i)$ has a_{i+1} as a suffix for $1 \leq i < m$, and $\tau(a_m)$ has a_1 as a suffix.

PROPOSITION 4. Suppose that τ has a generating fixed point $\mathbf{u} = u_0 u_1 u_2 \dots$. Then $a\mathbf{u} \in X_\tau$ if and only if a is part of a suffix cycle $a_1, a_2, \dots, a_m$, and $a_i u_0 \in \mathcal{L}_\tau$ for some $i \in \{1, \dots, m\}$.

Proof: If for some i , $a_i u_0 \in \mathcal{L}_\tau$, then since τ is primitive, there exists an n such that $a_i u_0 \in \tau^n(u_0)$, so $a_{i+1} \tau(u_0) \in \tau^{n+1}(u_0)$, and $a_i \tau^{nm}(u_0) \in \tau^{nm}(u_0)$ for each $n \geq 1$. It follows that $a_i \mathbf{u} \in X_\tau$, and that this is true for each $a_j \in \{a_1, \dots, a_m\}$.

Conversely, suppose that $a\mathbf{u} \in X_\tau$. We claim first that a is the suffix of some substitution word. If $a = \tau(b)$ for some b , we are done. If not, by Corollary 3, $a\mathbf{u} = \mathbf{w} \tau(\sigma \mathbf{x})$ where $\mathbf{w}$ is the suffix of some substitution word $\tau(x_0)$, and $\sigma \mathbf{x}$ is unique. Thus $\tau(\mathbf{u}) = \mathbf{u} = \sigma(\mathbf{w}) \tau(\sigma \mathbf{x})$, and by Corollary 3 again, $\sigma(\mathbf{w}) = \tau(u_0 \dots u_r)$ for some $r \geq 0$, so that a is the suffix of a substitution word.

Since (X_τ, σ) is minimal, σ is onto, which means that for each n there are words $\mathbf{w}_n$ of length n such that $\mathbf{w}_n a \mathbf{u} \in X_\tau$. Given N , choose n so that $n > |\tau^N(\alpha)|$ for all α . Thus $\mathbf{x} = \mathbf{w}_n a \mathbf{u} = \mathbf{w}' \tau^N(\sigma(\mathbf{y}))$ where $\mathbf{w}'$ is the suffix of $\tau^N(y_0)$, and Corollary 3 implies that for some $p > 0$, $\sigma^{p+1}(\mathbf{y}) = \mathbf{u}$ and $\mathbf{w}_n a = \mathbf{w}' \tau^N(\mathbf{y}_{[1,p]})$, so that a is the suffix of a τ^N word. Taking a subsequence, we conclude that there is some b such that a is the suffix of $\tau^N(b)$ for infinitely many N 's. Since the sequence of suffixes of $\tau^N(b)$ is eventually periodic, this implies that a is part of a suffix cycle. ■

It follows that if τ is left proper and suffix permutative, then $\mathbf{u}$ has $|\mathcal{A}|$ σ -preimages, and if τ is left and right proper, then $\mathbf{u}$ has one σ -preimage.

Let $\mathcal{W}_n := \{\mathbf{w} : \mathbf{w} = \sigma^k(\tau^n(a)) : a \in \mathcal{A}, 0 \leq k < |\tau^n(a)|\}$. If $\mathbf{w} \in \mathcal{W}_n$ and $\mathbf{w} = \sigma^k(\tau^n(a))$ with $k \geq 1$, let $r(\mathbf{w}) := |\{\alpha : \alpha\mathbf{w} \in \mathcal{W}_n\}|$, and let $\sigma^{-1}\mathbf{w} = \{\alpha\mathbf{w} : \alpha\mathbf{w} \in \mathcal{W}_n\}$.

PROPOSITION 5. *Suppose that $\mathbf{y}$ is not τ -fixed, and $\mathbf{y} \in [\mathbf{w}]$ with $\mathbf{w} \in \mathcal{W}_n$. Then $\mathbf{y}$ is a branch point only if $r(\mathbf{w}) > 1$. Conversely, suppose that there exists a sequence of words $\mathbf{w}_n \in \mathcal{W}_n$ with $|\mathbf{w}_n| \rightarrow \infty$, with $\cap_n [\mathbf{w}_n] = \mathbf{y}$. If $r(\mathbf{w}_n) > 1$ for each n , then $\mathbf{y}$ is a branch point.*

Proof: Since $\mathbf{y}$ is not fixed, then for some N , $\mathbf{y} \notin \tau^n(X_\tau)$ for each $n \geq N$, so that $\mathbf{y} = \mathbf{w}_n\tau(x)$ with $\mathbf{w}_n \in \mathcal{W}_n$. Then $\mathbf{y}$ is a branch point only if $r(\mathbf{w}_n) > 1$ for each $n \geq N$.

If $\mathbf{y} = \cap_n [\mathbf{w}_n]$ where $r(\mathbf{w}_n) > 1$ for each n , then passing down to a subsequence if necessary, we can assume that there are distinct α and β so that $\alpha\mathbf{w}_n$ and $\beta\mathbf{w}_n$ are elements of $\mathcal{W}_n$, hence $\mathcal{L}_\tau$. Thus $\alpha\mathbf{y}$ and $\beta\mathbf{y}$ are both elements of X_τ , so that $\mathbf{y}$ is a branch point. ■

Proposition 5 suggests an algorithm for generating all non- σ invertible points, and checking whether they are the τ -fixed point. We develop this algorithm in what follows, as our aim is to find the branch points of τ . If τ is defined on $\mathcal{A} = \{a_1, \dots, a_n\}$, pick $A_1 \subset \mathcal{A}$, where $\mathbf{s}_1$, the maximal proper common suffix of $\{\tau(a)\}_{a \in A_1}$, is non-empty. Thus for each a in A_1 , $\tau(a) = \mathbf{w}_a x_a \mathbf{s}_1$, and the set $A_2 := \{x_a : a \in A_1\}$ has at least two distinct elements. Call the set A_2 the set of $\mathbf{s}_1$ -predecessors in $\{\tau(a) : a \in A_1\}$. If $\{\tau(a) : a \in A_2\}$ has $\mathbf{s}_2$ as a maximal proper common suffix, let A_3 be the set of $\mathbf{s}_2$ -predecessors in $\{\tau(a) : a \in A_2\}$. Inductively, if A_k is the set of $\mathbf{s}_{k-1}$ -predecessors in $\{\tau(a) : a \in A_{k-1}\}$, let $\mathbf{s}_k$ be the maximal proper common suffix of $\{\tau(a) : a \in A_k\}$, and A_{k+1} the set of $\mathbf{s}_k$ -predecessors in $\{\tau(a) : a \in A_k\}$. This process may end after a finite number of steps, ie $A_k = \emptyset$ for some k - in which case we discard this A_1 and start over with another subset of $\mathcal{A}$. The sequence (A_k) is eventually periodic. For, the cardinality $|A_k|$ of the sets A_k cannot increase. If $|A_k|$ stays constant as k increases, then for some k and m , $A_k = A_{k+m}$, and so $\mathbf{s}_k = \mathbf{s}_{k+m}$, which in turn implies that $A_{k+1} = A_{k+m+1}$, and by induction that A_k is eventually periodic. If $|A_k|$ decreases, then repeat this argument. Thus $(\mathbf{s}_k)_{k \in \mathbb{N}}$ is also eventually periodic. Note that there can be more than one choice for A_1 which leads to a non-trivial sequence of $\mathbf{s}'_k$'s and A_k 's, and that there are only finitely many sequences $(\mathbf{s}_k)_{k \in \mathbb{N}}$ that can be generated, as A_1 ranges over all subsets of $\mathcal{A}$.

We illustrate this terminology with the following examples.

1. If $\tau(a) = aab$, and $\tau(b) = abb$, if $A_1 = \mathcal{A}$, then $A_n = \mathcal{A}$ and $\mathbf{s}_n = b$ for all $k \geq 2$. (In fact if τ is any substitution on a two letter alphabet whose substitution words have a proper maximal non-empty suffix, then it is always the case that $(\mathbf{s}_k)_{k \geq 1}$ is constant.)
2. If $\tau(a) = acb$, $\tau(b) = aba$ and $\tau(c) = aaa$, then the only possible choice for A_1 is $A_1 = \{b, c\}$. Here $\mathbf{s}_1 = a$, $A_2 = \{a, b\}$, and $\mathbf{s}_2 = \emptyset$ as $\tau(a)$ and $\tau(b)$ have no common suffix. Thus all choices of A_1 lead to sequences $(\mathbf{s}_n)$ which are eventually empty.
3. If $\tau(a) = acb$, $\tau(b) = aba$ and $\tau(c) = aca$, letting $A_1 = \{b, c\}$, then $A_k = A_1$ and $\mathbf{s}_k = \mathbf{s}_2$ for all $k \geq 2$.

LEMMA 6. *Let τ be defined on $\mathcal{A} = \{a_1, \dots, a_n\}$, and, given $A_1 \subset \mathcal{A}$, $(\mathbf{s}_m)_{m=1}^\infty$ and $(A_m)_{m=1}^\infty$ as defined and not eventually empty. Suppose that $\mathbf{s}_{k+p} = \mathbf{s}_k$ for $k \geq M$. Then*

1. *For $1 \leq n \leq M + p - 1$, the τ^n -substitution words have a maximal proper common suffix $\mathbf{S}_n = \left(\prod_{k=0}^{n-1} \tau^k(\mathbf{s}_{n-k})\right)$, and for $n = M + kp + j$, where $k \geq 1$ and $0 \leq j \leq p - 1$, the τ^n -substitution words have a maximal proper common suffix $\mathbf{S}_n = \mathbf{P}_n \mathbf{I}_n$, where*

$$\mathbf{P}_n = \left(\prod_{m=0}^j \tau^m(\mathbf{s}_{M+j-m}) \right) \prod_{l=0}^{k-1} \left(\prod_{i=0}^{p-1} \tau^{n-(M+(k-1-l)p+p-1-i)}(\mathbf{s}_{M+(p-1-i)}) \right) \quad (5)$$

and

$$\mathbf{I}_n = \left(\prod_{m=1}^{M-1} \tau^{n-(M-m)}(\mathbf{s}_{M-m}) \right). \quad (6)$$

2. If $(\mathbf{s}_n)$ is eventually the k -periodic sequence $(\mathbf{w}_i)_{i=1}^k$, then it defines k branch points, with each branch point $\mathbf{y}$ satisfying an equation of the form

$$(\mathbf{w}_i \tau(\mathbf{w}_{i+1}) \tau^2(\mathbf{w}_{i+2}) \dots \tau^{k-1}(\mathbf{w}_{i-1})) \tau(\mathbf{y}) = \mathbf{y}.$$

Also any branch point $\mathbf{y}$ which is non- τ -fixed arises in this fashion.

3. The substitution τ is quasi-invertible, with branch point $\mathbf{y}$ non τ -fixed, if and only if all τ -fixed points are invertible, and there is an $\mathbf{s} \in \mathcal{A}^+$ such that any $A_1 \subset \mathcal{A}$ generates a sequence $(\mathbf{s}_n)$ which is either eventually empty, or eventually the constant sequence $(\mathbf{s})$.
4. The substitution τ is quasi-invertible, with branch point $\mathbf{u}$, a τ -fixed point, if and only if $\mathbf{u}$ is the only τ -fixed branch point, and for all subsets $A_1 \subset \mathcal{A}$ that generate a sequence $(\mathbf{s}_n)$ which is not eventually empty, there is some N such that for $n > N$, there is a sequence $(k_n)_{n \geq N}$ of natural numbers such that the maximal common proper suffix of the $\{\tau^n(a) : a \in A_1\}$ have $\tau^{k_n}(\mathbf{u}_0)$ as a prefix, where $k_n \rightarrow \infty$.

Proof:

1. The proof of Equations 5 and 6 follow from the definitions of $(\mathbf{s}_k)$, and induction.
2. Note that if $\lim_{k \rightarrow \infty} \mathbf{S}_{n_k}$ exists, and $|\mathbf{P}_{n_k}| \rightarrow \infty$, then $\lim_{k \rightarrow \infty} \mathbf{S}_{n_k} = \lim_{k \rightarrow \infty} \mathbf{P}_{n_k}$, so we will assume that any sequence $(\mathbf{s}_k)$ is periodic.

In this case Equation 5 reduces to

$$\mathbf{S}_{np} = \mathbf{P}_{np} = \left(\prod_{m=0}^{p-1} \tau^m \mathbf{s}_{p-m} \right) \prod_{l=0}^{n-1} \left(\prod_{i=0}^{p-1} \tau^{lp+i}(\mathbf{s}_{p-i}) \right) \quad (7)$$

Here $\mathbf{S}_{(n+1)p}$ is a prefix of $\mathbf{S}_{np}$ for $n \geq 1$, and if $|\mathbf{S}_{np}| \rightarrow \infty$, then $\mathbf{y} := \lim_{n \rightarrow \infty} \mathbf{S}_{np}$ exists, and by Proposition 5 has at least 2 σ -preimages. Note that $\mathbf{y}$ satisfies the equation

$$\left(\prod_{m=0}^{p-1} \tau^m \mathbf{s}_{p-m} \right) \tau^p(\mathbf{y}) = \mathbf{y}. \quad (8)$$

A similar argument applied to the sequence $(\mathbf{S}_{(n+1)p+i})_n$ yields the other branch points. Conversely, if $\mathbf{y}$ is a non- τ -fixed k -branch point, then there are k letters $\{a_1^n, a_2^n, \dots, a_k^n\}$ whose τ^n words share a maximal common proper suffix $\mathbf{T}_n$ and such that $\mathbf{y} \in [\mathbf{T}_n]$. Moving to a subsequence if necessary, we can assume that these k letters are independent of n and so if $A_1 = \{a_1, \dots, a_k\}$, the sequence $(\mathbf{T}_n)$ is identical to the sequence $(\mathbf{S}_n)$ generated by A_1 . The result follows.

3. Statements 3 and 4 now follow from Part 2. ■

Examples

4. (Example 1) If $\tau(a) = aab$, and $\tau(b) = abb$, then the unique branch point $\mathbf{y}$ satisfies $b\tau(\mathbf{y}) = \mathbf{y}$.
5. (Example 2) If $\tau(a) = acb$, $\tau(b) = aba$ and $\tau(c) = aaa$, then only $\mathbf{u}$ is a 2-branch point.
6. (Example 3) If $\tau(a) = acb$, $\tau(b) = aba$ and $\tau(c) = aca$, then $\{a, b\}$ is a suffix cycle, so $\mathbf{u}$ is a 2-branch point. Letting $A_1 = \{b, c\}$, we obtain the (only other) 2-branch point $\mathbf{y}$ satisfying $a\tau(\mathbf{y}) = \mathbf{y}$, and $\mathbf{y} \neq \mathbf{u}$.
7. If $\tau(a) = abc$, $\tau(b) = aacc$, and $\tau(c) = abcc$, then $\mathbf{u}$ is σ -invertible, and if $A_1 = \{b, c\}$, $\mathbf{s}_{2k+1} = cc$ and $\mathbf{s}_{2k} = c$ for $k \geq 1$, and no other choice of A_1 generates any other eventually different sequence $(\mathbf{s}_k)$. Thus (X_τ, σ) has two branch points, $\mathbf{y}$ and $\mathbf{y}^*$, satisfying $\mathbf{y} = c\tau(cc)\tau(\mathbf{y})$ and $\mathbf{y}^* = cc\tau(c)\tau(\mathbf{y}^*)$, and these two points are distinct.

8. If $\tau(a) = bbad$, $\tau(b) = ab$, $\tau(c) = ad$, and $\tau(d) = dac$, then τ has 3 fixed points, with $\tau^\infty(a)$ a 3-branch point, $\tau^\infty(b)$ a 2-branch point, and where $\tau^\infty(d)$ is σ -invertible. If $A_1 = \{a, b\}$ then the words $\{\tau^n(a), \tau^n(b)\}$ have $\tau^n(b)$ as a maximal common suffix, and these words also converge to u .

We remark that the notion of a branch point has been studied, for example in [BDH03]. There the authors work with two dimensional sequences and define the notion of a *right asymptotic* point. The relationship is that if x is a one-sided k -branch point, then this leads to k right asymptotic two sided points. The authors find upper bounds on the number of right asymptotic points - being $2n - 1$ if $|\mathcal{A}| = n$ and τ is left proper, and $2(n - l) + rl$ in general, if τ has l proper prefixes and r proper suffixes (Proposition 9 and Theorem 1 of [BDH03]), answering a question in [HZ01]. We digress here to show that if $2 \leq i \leq 2n - 1$, there exists a left proper τ which has i right asymptotic points. A similar argument will work to create non-proper substitutions having a prescribed (appropriately constrained) number of branch points.

PROPOSITION 7. *Let τ be left proper on n letters. For any m -tuples of natural numbers $(k_1, k_2, \dots, k_m)$, $(j_1, j_2, \dots, j_m)$ where*

1. $2 \leq k_1 < k_2 < \dots < k_m \leq n$
2. $k_1 + j_1(k_1 - 1) \leq n + 1$, and for each $i \geq 2$, $k_i + j_i(k_i - 1) \leq n + 1 - \sum_{l=1}^{i-1} k_l + j_l(k_l - 1)$
and,
3. $\sum_{l=1}^m k_l + j_l(k_l - 1) \leq n + m$,

there exists a left proper τ on n letters, with j_i k_i -branch points, for $1 \leq i \leq m$, and no other branch points.

Proof: We first prove the special case $m = 1$, $k_1 = 2$, and $j_1 \leq n - 1$, which would generate a substitution with j_1 2-branch points, and so $2j_1$ left asymptotic points. Letting $\mathcal{A} = \{x_1, \dots, x_n\}$, choose some letter a so that all τ -substitution words start with a . Let $\tau(x_1)$ have x_1 as suffix and let all other $\tau(x_i)$'s have x_2 as suffix. As long as we ensure later that both $x_1a, x_2a \in \mathcal{L}_\tau$, then the τ -fixed point is a 2-branch point. We define $\tau(x_i)$ by successively adding letters from the right hand side, moving towards the left. Define the penultimate letter of $\tau(x_2)$ to be x_2 , and the penultimate letter of all other $\tau(x_i)$ to be x_3 . Then we have a second branch point (satisfying the equation $x_2\tau(y) = \tau(y)$). At the l th stage (for $l \leq n - 1$), fill in the l th column from the right of $\tau(x_i)$ by putting down an x_l for $\tau(x_l)$, and x_{l+1} elsewhere. In this way you get $n - 1$ 2-branch points, $n - 2$ of them satisfying $x_l(\tau(y)) = \tau(y)$ for $2 \leq l \leq n - 1$ (and the last τ -fixed). If $j_1 < n - 1$, you stop the above procedure after having defined the j_1 -st column from the right. To ensure you get no more branch points (and also that τ is injective, if required) start filling in columns with x_1 and x_2 (or x_i and x_{i+1} , where $i \leq j_1$). Then any new branch points satisfy an already defined equation - $x_i\tau(y) = y$ for some $i \leq j_1$.

In fact the above special case where $j_1 = n - 1$ is how one maximises the number of branch, and left asymptotic, points. These 2-branch points can be packed in an optimal way. To obtain j k -branch points, one needs $k + k(j - 1)$ letters, replicating the above example, but at each stage when defining the τ substitution words, having k different letters in each column that you define. To ensure the branch points that you create are distinct, the letters should change from column to column. When the desired number of branch points have been obtained, start filling in new columns with combinations of letters that have already been used to create a previous branch point (for example use a subset of $x_1, x_2, \dots, x_k$ if they were used to create a k -branch point).

■

Unilateral recognizability is less amenable than bilateral recognizability, which all aperiodic primitive substitutions possess, and for whom the sequence of partitions $(\mathcal{P}_n)$ defined by

$$\mathcal{P}_n := \{[\sigma^k(\tau^n(a))] : a \in \mathcal{A}, 0 \leq k < |\tau^n(a)|\}$$

is a nested sequence which spans the topology of $(X_\tau^\mathbb{Z}, \sigma)$ (see Proposition 14, [DHS99]). In the one sided subshift (X_τ, σ) , the sets in $\mathcal{P}_n$ need not be disjoint. For example, if τ has a branch point $\mathbf{y} \neq \mathbf{u}$, then, for each n , there are letters a, b , and numbers n_1 and n_2 such that $|\tau^n(a)| - n_1 = |\tau^n(b)| - n_2$ and $\sigma^{n_1}(\tau^n(a)) = \sigma^{n_2}(\tau^n(b))$, while $\sigma^{n_1-1}(\tau^n(a)) \neq \sigma^{n_2-1}(\tau^n(b))$. If τ is such that all branch points are fixed in X_τ , this does not happen.

PROPOSITION 8. *Suppose that the k branch points $\{\mathbf{u}_1, \dots, \mathbf{u}_k\}$ of τ are τ -fixed.*

Then

1. *For every $n > 0$, $\mathcal{P}_n$ is a clopen partition, and the sequence of bases is decreasing.*
2. *For each n , $\mathcal{P}_{n+1}$ is a refinement of $\mathcal{P}_n$: i.e. every element of $\mathcal{P}_{n+1}$ is contained in an element of $\mathcal{P}_n$.*
3. *The intersection of the bases of $\mathcal{P}_n$ is $\{\mathbf{u}_1, \dots, \mathbf{u}_k\}$, and the sequence $(\mathcal{P}_n)$ spans the topology of (X_τ, σ) .*

Proof:

1. We show that $\mathcal{P}_1$ is clopen. First suppose that $(\mathbf{y}_n)$ is a sequence in $\tau[a]$ and $\mathbf{y}_n \rightarrow \mathbf{y}$. Since $\tau(X_\tau)$ is closed, by Proposition 2, then $\mathbf{y} \in \tau(X_\tau)$. If $\mathbf{y}_n = \tau(\mathbf{x}_n)$ and $\mathbf{x}$ is a limit point of $(\mathbf{x}_n)$, then $\mathbf{x} \in [a]$ and $\mathbf{y} = \tau(\mathbf{x}) \in \tau[a]$. Using Corollary 3, the sets $\{\tau[a] : a \in \mathcal{A}\}$ are disjoint, and their union, $\tau(X_\tau)$, is a clopen set, so they are also. If $0 < k < |\tau(a)|$ and $(\mathbf{y}_n)$ is a sequence of elements in $[\sigma^k(\tau(a))]$, then $\mathbf{y}_n = \sigma^k(\mathbf{x}_n)$ where $\mathbf{x}_n \in \tau[a]$. If $\mathbf{x}$ is a limit point of $(\mathbf{x}_n)$ then it must also be in $\tau[a]$, and $\mathbf{y} = \sigma^k(\mathbf{x}) \in \sigma^k(\tau[a])$. So the sets $\sigma^k(\tau[a])$ are closed, and if the branch points are all τ -fixed, then these sets are also disjoint, and so clopen.
2. Take an element $\mathbf{p} \in \mathcal{P}_{n+1}$: we need to show that $[\mathbf{p}] \subset [\mathbf{p}']$ where $[\mathbf{p}'] \in \mathcal{P}_n$. Suppose that $\mathbf{p} = \sigma^k(\tau^{n+1}(a))$, where $0 < k < |\tau^{n+1}(a)|$. As in the proof of Proposition 14, in [DHS99], write $\tau^{n+1}(a) = \tau^n(a_1) \tau^n(a_2) \dots \tau^n(a_m)$ where $\tau(a) = a_1 a_2 \dots a_m$. For some j , $|\tau^n(a_1 \dots a_j)| \leq k < |\tau^n(a_1 \dots a_{j+1})|$, so let $l := k - |\tau^n(a_1 \dots a_j)|$. Then $\sigma^k(\tau^{n+1}([a])) \subset \sigma^l \tau^n[a']$, where $a' = a_{j+1}$. Let $\mathbf{p}' := \sigma^l \tau^n[a']$.
3. To see that $(\mathcal{P}_n)$ span the topology of (X_τ, σ) , we imitate the proof of Proposition 14, [DHS99]. Given m positive, we show that for all n large, each element $\mathbf{p}$ of $\mathcal{P}_n$ is contained in some $\mathbf{c}$ where $\mathbf{c}$ is a word of length $m+1$. Suppose that all τ -substitution words start with l . Write $l_n = |\tau^{n-1}(l)|$. Choose n so that $l_n > m$ (minimality implies that $\lim_{n \rightarrow \infty} l_n = \infty$).

Fix $\mathbf{p} \in \mathcal{P}_n$, and suppose that $\mathbf{p} = \sigma^k(\tau^n(a))$ for some $a \in \mathcal{A}$ and $k \in [0, |\tau^n(a)|]$. If $\mathbf{y} \in [\mathbf{p}]$ then there is some $\mathbf{x} \in [a]$ with $\mathbf{y} = \sigma^k \tau^n(\mathbf{x})$. Since τ is left proper, $\tau^n(a) \tau^{n-1}(l)$ is a prefix of $\tau^n(\mathbf{x})$, and so $\mathbf{y} \in [\sigma^k \tau^n(a) \tau^{n-1}(l)]$, and this last cylinder set has length greater than m . So, $\mathbf{y}$ starts with a block of length m depending only on $\mathbf{p}$ and not on $\mathbf{y}$. ■

4. Bratteli Diagrams

A *stationary Bratteli diagram* $\mathcal{B} = (\mathcal{V}, \mathcal{E})$ is an infinite directed graph with *vertex set* $\mathcal{V} = \bigsqcup_{n=0}^{\infty} V_n$ and *edge set* $\mathcal{E} = \bigsqcup_{n=1}^{\infty} E_n$, where $V_n = V$ for each $n \geq 1$, $E_n = E$ for each $n \geq 1$, $V_0 = \{v_0\}$, and for each $n \geq 1$ there is a *source map* $s : E_n \rightarrow V_{n-1}$ and a *range map* $r : E_n \rightarrow V_{n+1}$. If $|V_n| = d$, the *incidence matrix* $F = (f_{i,j})_{1 \leq i, j \leq d}$ is defined

by $f_{i,j} := |\{x \in E_n : s(x) = j, r(x) = i\}|$ for any $n \geq 2$. If $n = 1$, the incidence matrix F_1 describing E_1 is a $d \times 1$ vector defined in a similar way. We assume that $s^{-1}(v) \neq \emptyset$ for each $v \in \mathcal{V}$ and $r^{-1}(v) \neq \emptyset$ for all $v \in \mathcal{V} \setminus v_0$. We will use $x, y, \dots$ when referring to edges, and $a, b, \dots$ when referring to vertices. We say that $\mathcal{B}$ is *simple* if some power of F has all positive entries.

A finite set of edges $\{x_{n+k}\}_{k=1}^K$, with $s(x_{n+k+1}) = r(x_{n+k})$ for $1 \leq k \leq K-1$, is called a *path* from $s(x_{n+1})$ to $r(x_{n+K})$; we can extend the maps s and r to paths by defining $s(\{x_{n+k}\}_{k=1}^K) = s(x_{n+1})$ and $r(\{x_{n+k}\}_{k=1}^K) = r(x_{n+K})$. Similarly an *infinite path* in $\mathcal{B}$ is a sequence $\mathbf{x} = (x_n)_{n=1}^\infty$, with $x_n \in E_n$ for $n \geq 1$, and $s(x_{n+1}) = r(x_n)$ for $n \geq 1$. The set of all infinite paths in $\mathcal{B}$ will be denoted $X_{\mathcal{B}}$ (a subset of $\prod_{n \geq 1} E_n$), and $X_{\mathcal{B}}$ is endowed with the topology induced from the product topology on $\prod_{n \geq 1} E_n$, so that $X_{\mathcal{B}}$ is a compact metric space. If $\mathbf{x} = (x_n)_{n=1}^\infty$ and $\mathbf{x}' = (x'_n)_{n=1}^\infty$ are two elements in $X_{\mathcal{B}}$, we write $\mathbf{x} \sim \mathbf{x}'$ if there is some N such that $x_n = x'_n$ for all $n \geq N$. It follows that $\sim$ is an equivalence relation (the *tail equivalence relation*), and if $\mathcal{B}$ is simple, then each equivalence class for $\sim$ is dense in $X_{\mathcal{B}}$, and $X_{\mathcal{B}}$ has no isolated points, making it a Cantor space.

We will sometimes *telescope* $\mathcal{B}$ to $\mathcal{B}'$ - for our purposes we need only define a restricted version of this procedure. Let k be given; then $\mathcal{B}^{(k)} = (\mathcal{V}^{(k)}, \mathcal{E}^{(k)})$ is a *telescoping* of $\mathcal{B} = (\mathcal{V}, \mathcal{E})$ if $V_n^{(k)} = V_{nk}$ (with the vertex $v \in V_{nk}$ labelled as $v \in V_n^{(k)}$), and $F^{(k)} := F^k$, so that the number of edges from $v_n \in V_n^{(k)}$ to $v_{n+1} \in V_{n+1}^{(k)}$ is the number of paths from $v_n \in V_{nk}$ to $v_{n+1} \in V_{(n+1)k}$. Conversely, we can perform a *splitting* by introducing a new level between two consecutive levels V_{n-1} and V_n , and new edge sets, so that if the resulting stationary Bratteli diagram is $\mathcal{B}' = (\mathcal{V}', \mathcal{E}')$, and we telescope to every other level, we obtain the original $\mathcal{B} = (\mathcal{V}, \mathcal{E})$. We will split diagrams only when a vertex in V_1 has more than one edge connecting it to v_0 . We consider two Bratteli diagrams $\mathcal{B}$ and $\mathcal{B}'$ *equivalent* if $\mathcal{B}'$ can be obtained from $\mathcal{B}$ by relabeling, telescoping and splitting, so that when we talk about a Bratteli diagram we are talking about an equivalence class of diagrams.

4.0.1. Ordering $X_{\mathcal{B}}$ For each $n \geq 1$ and each $a \in V_n$, we put a linear order on $\{x \in E_n : r(x) = a\}$. In this article we assume that the order we put on $\{x \in E_n : r(x) = a\}$ depends only on a and not n if $n > 1$. We also call this ordering *stationary*. Note that we do not put any conditions on the ordering of E_1 . Elements of $E_n(a)$ will then be labelled $0, 1, 2, \dots$ according to their order, and we say $\mathcal{B}$ is *ordered*. We will also write $x < y$ if x and y have the same range and x has a smaller labelling than y . Every stationary ordering is generated by a substitution τ on $\mathcal{A}$, an alphabet of size $|V|$: if the i th incoming edge to a has source b , then define the i th letter of $\tau(a)$ to be b . Conversely to each substitution τ on $\mathcal{A}$, we can define the (stationary, ordered) *Bratteli diagram associated with τ* , denoted $\mathcal{B}_\tau$: there is exactly one edge from each vertex in V_1 to v_0 , let the transition matrix for τ be the incidence matrix F for $\mathcal{B}$, and order $\mathcal{B}$ using the order of appearance of letters in the substitution words: if the i -th letter of $\tau(a)$ is b , then give one edge x with $s(x) = b$ and $r(x) = a$ the ordering i . If $\mathcal{B}$ is the diagram defined by the transition matrix M for τ then $\mathcal{B}^{(N)}$ is the diagram defined by the transition matrix M^N for τ^N . We will use the substitution τ to denote the ordering on $\mathcal{B}$ and write $\mathcal{B}_\tau$ for the ordered diagram - in this case $\mathcal{B}_{\tau^N}^{(N)}$ is the ordered diagram associated to τ^N .

For $m < n$, the order τ induces a partial ordering on paths from V_m to V_n : the two paths $\mathbf{x} = x_{m+1} \dots x_n$ and $\mathbf{x}' = x'_{m+1} \dots x'_n$ with source in E_m and common range in E_n are comparable, with $\mathbf{x} < \mathbf{x}'$ if there is some $k \in [m+1, n]$ with $x_k < x'_k$ and $x_j = x'_j$ for $k+1 \leq j \leq n$. If $1 \leq m < n$, then the number of paths in $\mathcal{B}$ from $b \in \mathcal{V}_m$ to $a \in \mathcal{V}_n$ is the number of occurrences of b in $\tau^{(n-m)}(a)$. If τ is primitive then there is a positive k such that for any two letters a and b , there is at least one path from $b \in \mathcal{V}_n$ to $a \in \mathcal{V}_{n+k}$.

The ordering τ on $\mathcal{B}$ extends to an order on tail equivalence classes in $X_{\mathcal{B}}$: if $\mathbf{x} \sim \mathbf{x}'$, we write $\mathbf{x} < \mathbf{x}'$ if an initial segment $x_1 \dots x_N < x'_1 \dots x'_N$. An infinite path is *maximal* (*minimal*) if all the edges making up the path are maximal (minimal). For any τ there is a power τ^N such that all substitution words of τ^N have prefixes which generate right infinite τ -fixed points, and suffixes which generate left infinite τ -fixed points. Each one of these prefixes generates a distinct minimal path in $X_{\mathcal{B}}$, and each such suffix generates a distinct maximal path in $X_{\mathcal{B}}$. If $\{v_1, v_2, \dots, v_n\}$ ($\{w_1, w_2, \dots, w_k\}$) are these suffixes (prefixes), we call these the vertices that determine the maximal (minimal) paths in $X_{\mathcal{B}}$. Note that the orderings τ and τ^N generate the same number of maximal and minimal paths in $X_{\mathcal{B}}$ and $X_{\mathcal{B}^{(N)}}$ respectively. If τ generates a minimal subshift, then $k \leq n$. Note that all maximal and minimal paths in $X_{\mathcal{B}}$ are (shift) periodic, and those in $X_{\mathcal{B}^{(N)}}$ are (shift) fixed. Let $X_{\max} := \{\mathbf{M}_1, \mathbf{M}_2, \dots, \mathbf{M}_n\}$ and $X_{\min} = \{\mathbf{m}_1, \mathbf{m}_2, \dots, \mathbf{m}_k\}$ denote the sets of maximal and minimal paths respectively in $X_{\mathcal{B}}$. They are both non-empty.

If $\mathbf{x} = (x_n)_{n=1}^{\infty}$ is not maximal, let k be the smallest integer such that x_k is not a maximal edge, and let y_k be the successor of x_k in $\{x : r(x) = r(x_k)\}$. Then the *successor* $V_{\tau}(\mathbf{x})$ is defined to be $V_{\tau}(\mathbf{x}) = 0 \dots 0 y_k x_{k+1} x_{k+2} \dots$, where $0 \dots 0$ is the unique minimal path ending at the source of y_k . Similarly, every non-minimal path has a unique *predecessor*. Simple ordered Bratteli diagrams which have a unique minimal and maximal element are called *proper*, and those with a unique minimal element $\mathbf{m}$ are called *semi-proper*. If $\mathcal{B}_{\tau}$ is proper, then V_{τ} is a homeomorphism, and if $\mathcal{B}_{\tau}$ is semi-proper, then V_{τ} can be extended to a continuous surjection on $X_{\mathcal{B}}$ by setting $V_{\tau}(\mathbf{M}_i) = \mathbf{m}$ for all maximal elements $\mathbf{M}_i$. If $\mathcal{B}_{\tau}$ is not semi-proper, the continuity of V_{τ} turns out to depend entirely on $\mathcal{L}_{\tau}$: we address this in the next proposition. We remark that a special case of this result is implicit in the formula on Line 12, Page 5 of [HZ01], when the number of maximal elements equals the number of minimal elements in $\mathcal{B}_{\tau}$.

PROPOSITION 9. *Suppose that $\mathcal{B} = \mathcal{B}_{\tau}$ is a stationary ordered Bratteli diagram, which has maximal elements $\{\mathbf{M}_i\}_{i=1}^n$ and minimal elements $\{\mathbf{m}_i\}_{i=1}^k$ where $k \leq n$, determined by the vertices $\{v_1, v_2, \dots, v_n\}$ and $\{w_1, w_2, \dots, w_k\}$ respectively. Then V_{τ} can be extended to a continuous mapping $V_{\tau} : X_{\mathcal{B}} \rightarrow X_{\mathcal{B}}$ if and only if for each $v_i \in \{v_1, v_2, \dots, v_n\}$ there exists a unique $w_j \in \{w_1, w_2, \dots, w_k\}$ such that $v_i w_j \in \mathcal{L}_{\tau}$.*

Proof: We first prove the result for substitutions which generate maximal and minimal paths in $X_{\mathcal{B}}$ that are σ -fixed. In other words, suppose that all τ -substitution words have a prefix (suffix) which generates a right (left) infinite fixed point. Suppose also that all words of length two in $\mathcal{L}_{\tau}$ appear in the τ -substitution words.

First suppose that V_{τ} can be extended to a continuous map $V_{\tau} : X_{\mathcal{B}} \rightarrow X_{\mathcal{B}}$. If $f(\mathbf{M}_i) = \mathbf{m}_j$, then for large N , elements in the cylinder set $[M_i^N e]$, with e not a maximal edge, are mapped, using continuity, into the cylinder set $[m_j^{N-1}]$. If $s(e) = s(e+1) = v$ then $v_i w_j$ is a subword of $\tau^2(v)$, and so in $\mathcal{L}_{\tau}$. Suppose that $v_i w_{j'} \in \mathcal{L}_{\tau}$; we show that $j' = j$. Suppose that $v_i w_{j'}$ appears in a τ -substitution word $\tau(v)$, then there exist edges $\alpha, \alpha+1$ with sources v_i and $w_{j'}$ respectively and range v . For large N , take the cylinder set $[M_i^N \alpha]$; its image under $V_{\mathcal{B}}$ lies in $[m_{j'}^N \alpha + 1]$ and continuity means that $j = j'$.

Conversely suppose that for each i , there is a unique j with $v_i w_j \in \mathcal{L}_{\tau}$. Define $V_{\tau}(\mathbf{M}_i) := \mathbf{m}_j$. If V_{τ} is not continuous, then there are distinct edges α, β with source v_i , but for large N , $V_{\tau}(M_i^N \alpha) \in [w_{j'}^{N-1}]$ and $V_{\tau}(M_i^N \beta) \in [w_{j''}^{N-1}]$, where $j' \neq j''$, and one of j', j'' , say j' , is distinct from j . If $r(\alpha) = v$, then $v_i w_{j'} \in \tau^2(v)$, so that $v_i w_{j'} \in \mathcal{L}_{\tau}$, a contradiction.

For a general substitution τ , there is some power τ^N whose Bratteli diagram $\mathcal{B}^{(N)}$ satisfies all our initial assumptions. Also, $(X_{\mathcal{B}}, V_{\tau})$ is continuous if and only if $(X_{\mathcal{B}^{(N)}}, V_{\tau^N})$ is

continuous. Finally the substitutions τ and τ^N generate the same sets $\{v_1, v_2, \dots, v_n\}$ and $\{w_1, w_2, \dots, w_k\}$, and also $\mathcal{L}_\tau = \mathcal{L}_{\tau^N}$. The result follows. $\blacksquare$

Let us call a stationary ordered diagram $\mathcal{B}_\tau$ *closing* if it satisfies the conditions of Proposition 9. If $\mathcal{B}_\tau$ is closing, then $(X_\mathcal{B}, V_\tau)$ is called a *Bratteli-Vershik* or *adic* system. Since V_τ orbits are equivalence classes for $\sim$, if τ is primitive, so that $\mathcal{B}_\tau$ is simple, then every V_τ orbit is dense. Note that if there are multiple edges between v_0 and vertices in V_1 , this has no bearing on the continuity of V_τ : ie for any transition matrix F_1 and any order put on E_1 , the resulting map V_τ (we abuse notation here, as different F_1 's generate different V_τ 's) is continuous if and only if the V_τ is continuous for the choice $F_1 := (1, 1, \dots, 1)$.

The following is the specialised version of Theorem 4.6 in [HPS92] that we will need:

PROPOSITION 10. *Let (X_τ, σ) be a substitution subshift where all k branch points $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\}$ are τ -fixed. If V_τ is continuous, then $(X_\tau, \sigma) \stackrel{\Phi}{\cong} (X_\mathcal{B}, V_\tau)$, with the points $\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\}$ mapped to the k minimal elements of $X_\mathcal{B}$.*

Proof: This follows from Proposition 8, and standard constructions of Φ , as for example in the proof of Theorem 4.6 in [HPS92].

5. Quasi-invertible substitutions

In this section we obtain adic representations for any quasi-invertible substitution τ . Here we are in the situation where we do not have to worry about the continuity of V_τ , so that to apply Proposition 10 we need to find some τ' with $(X_\tau, \sigma) \cong (X_{\tau'}^\mathbb{N}, \sigma)$, and where the branch point of τ' is τ' -fixed.

LEMMA 11. *Suppose that τ is defined as $\tau(a_i) = \mathbf{p}_i x_i \mathbf{s}_1$, where $\{x_1, \dots, x_n\}$ has at least two distinct elements, and where if $A_1 = \mathcal{A}$, then $\mathbf{s}_k = \mathbf{s}_1$ for $k \geq 1$. Define the substitution τ^* as $\tau^*(a_i) = \mathbf{s}_1 \mathbf{p}_i x_i$.*

1. *For each $n \geq 0$, and each $\mathbf{w} \in \mathcal{L}_\tau$, $\tau^*(\tau^n(\mathbf{w}))\mathbf{s}_1 = \mathbf{s}_1 \tau^{n+1}(\mathbf{w})$.*
2. *If $\mathbf{S}_n$ is as in Lemma 6, then for each $n \geq 2$, $\mathbf{S}_n = \tau^*(\mathbf{S}_{n-1})\mathbf{s}_1$.*
3. *$\mathbf{S}_n \subset (\tau^*)^{n-1}(\mathbf{s}_1)$ for each $n > 1$.*

Proof:

1. Note that $\tau^*(a)\mathbf{s}_1 = \mathbf{s}_1 \mathbf{p}_a x_a \mathbf{s}_1 = \mathbf{s}_1 \tau(a)$ for any $a \in \mathcal{A}$. The general proof for longer words follows by concatenation.
2. First by Lemma 6 and Part 1 of this lemma, $\mathbf{S}_2 = \mathbf{s}_1 \tau(\mathbf{s}_1) = \tau^*(\mathbf{s}_1)\mathbf{s}_1$. Assuming that $\mathbf{S}_n = \tau^*(\mathbf{S}_{n-1})\mathbf{s}_1$, we have

$$\mathbf{S}_{n+1} = \mathbf{S}_n \tau^n(\mathbf{s}_1) \stackrel{IH}{=} \tau^*(\mathbf{S}_{n-1})\mathbf{s}_1 \tau^n(\mathbf{s}_1) \stackrel{Part1}{=} \tau^*(\mathbf{S}_{n-1})\tau^*(\tau^{n-1}(\mathbf{s}_1))\mathbf{s}_1 = \tau^*(\mathbf{S}_n)\mathbf{s}_1.$$

3. The case $n = 2$ is clear. Assuming $\mathbf{S}_n \subset (\tau^*)^{n-1}(\mathbf{s}_1)$,

$$\mathbf{S}_{n+1} \stackrel{Part2}{=} \tau^*(\mathbf{S}_n)\mathbf{s}_1 \stackrel{IH}{\subset} \tau^*((\tau^*)^{n-1}(\mathbf{s}_1))\mathbf{s}_1 = (\tau^*)^n(\mathbf{s}_1)\mathbf{s}_1 \subset (\tau^*)^n(\mathbf{s}_1).$$

$\blacksquare$

COROLLARY 12. *Let τ be left proper and quasi-invertible, with branch point $\mathbf{y} \neq \mathbf{u}$. Then some power of τ is right proper, and there exists a left proper, quasi-invertible substitution τ^* with $\tau^*(\mathbf{y}) = \mathbf{y}$, and $(X_\tau, \sigma) = (X_{\tau^*}, \sigma)$.*

Proof: Since the fixed point $\mathbf{u}$ is σ -invertible, Proposition 4 tells us that some τ (or some power of τ) is right proper. If $\mathbf{y}$ is the branch point for τ , then Lemma 6 tells us that there

exists an $\mathbf{s}$ such that if A is any subset of $\mathcal{A}$, it generates an eventually fixed sequence $(\mathbf{s}_k)$ where $\mathbf{s}_k = \mathbf{s}$ for $k \geq n_A$. Since there are finitely many such sequences we can assume, taking a power if necessary, that $n_A = 1$. Now Lemma 11 can be applied, taking τ^* as defined. The branch point is $\mathbf{y} := \lim_{n \rightarrow \infty} \mathbf{S}_n$; that it exists and has at least two σ -preimages follows from Lemma 6. That it is τ^* -fixed follows from Part 3 of Lemma 11. Thus $X_{\tau^*} \subset X_\tau$, and by minimality, this inclusion is an equality. ■

The connection between Bratteli-Vershik systems and one sided left proper substitutions is given by the next result. It is the appropriate generalization of [DHS99, Prop 20] to one sided substitution systems.

THEOREM 13. *If τ is left proper and quasi-invertible, with fixed point $\mathbf{u}$, there exists a semi-proper, stationary Bratteli diagram $\mathcal{B}$ which is semi-proper, and stationary, such that $(X_{\mathcal{B}}, V_{\mathcal{B}})$ is topologically conjugate to (X_τ, σ) .*

Proof: If $\mathbf{u}$ is the branch point, the sequence of partitions $\mathcal{P}_n$ defined in Section 3 are a refining sequence of partitions, by Proposition 8, which generate the topology of (X_τ, σ) . Proposition 10 tells us that $(X_\tau, \sigma) \cong (X_{\mathcal{B}}, V_\tau)$. If τ is left proper and the branch point $\mathbf{y} \neq \mathbf{u}$, then by Lemma 4, τ (or some power of τ) has to be right proper. Work with the substitution τ^* in Corollary 12. As above, (X_{τ^*}, σ) is conjugate to $(X_{\mathcal{B}^*}, V_{\tau^*})$, and since $(X_\tau, \sigma) = (X_{\tau^*}, \sigma)$, the result follows. ■

5.1. Return words and induced substitutions We now introduce concepts needed to extend Theorem 13 to non-left proper substitutions. Suppose that $T : X \rightarrow X$ is a minimal continuous transformation with X a Cantor space - henceforth called a Cantor system. If $U \subset X$ is a clopen set, the *system (U, T_U) induced by (X, T) on U* , is defined by $T_U(\mathbf{x}) = T^n(\mathbf{x})$, where n is the least positive natural number such that $T^n(\mathbf{x}) \in U$. T_U is well defined since T is minimal, and since X is compact, n can only take a finite number of values. The Cantor systems (X_1, T_1) and (X_2, T_2) are (topologically) *Kakutani equivalent* ([PS73] is first reference of this) if there exist clopen sets $U_i \subset X_i$ such that the respective induced systems are isomorphic. Topological Kakutani equivalence is much more stringent than measurable Kakutani equivalence - for example, all rank one transformations are measurably Kakutani equivalent, but it can be shown, for example, that the *Chacon* substitution $\tau(0) = 0010$, $\tau(1) = 1$ is Kakutani equivalent only to the substitutions $\tau_1(0) = 001^k0$, $\tau_1(1) = 1$, or $\tau_2(0) = 01^k00$, $\tau_2(1) = 1$.

Conversely, given an induced system (U, σ_U) , where (X, σ) is minimal, for some finite H , we can define the *height function* $h : U \rightarrow \{1, \dots, H\}$ given by $h(\mathbf{x}) = k$ if and only if $\sigma^i(\mathbf{x}) \notin U$ for $1 \leq i \leq k-1$ but $\sigma^k(\mathbf{x}) \in U$. For each k in $\{1, \dots, H\}$, and $0 \leq k-1$, let $U_k^i = \sigma^i(h^{-1}(\{k\}))$. Since (X, σ) is minimal, $X = \bigcup_{k=1}^H \bigcup_{j=0}^{k-1} U_k^j$.

LEMMA 14. *If (X_τ, σ) is quasi-invertible, with branch point in the clopen set U , then the sets $\{U_k^j : 1 \leq k \leq H, 0 \leq j \leq k-1\}$ form a clopen partition of X_τ .*

Proof: It is straightforward that each U_k^0 is clopen, by their definition. If $(\mathbf{y}_n) \subset U_k^i$ and $\mathbf{y}_n \rightarrow \mathbf{y}$, then for each n there is some $\mathbf{x}_n \in U_k^0$ with $\sigma^k(\mathbf{x}_n) = \mathbf{y}_n$. If $\mathbf{x}$ is a limit point of the points $\mathbf{x}_n$, then $\mathbf{x} \in U_k^0$ and $\mathbf{y} = \sigma^i(\mathbf{x})$. Thus U_k^i is closed. If the branch point in X_τ is in U , these sets are also disjoint. The result follows. ■

If we can partition X_τ as in Lemma 14, we say that (X_τ, σ) is a *primitive* of (U, σ_U) . If (U, T_U) has an appropriate adic representation, it is now possible to extend this representation to (X_τ, T) , as in the proof of Theorem 3.8, in [GPS95]:

THEOREM 15. *Let $(X_{\mathcal{B}}, V_{\mathcal{B}})$ be an adic system, where $\mathcal{B}$ is semi proper, simple, and with M maximal elements. Suppose that the aperiodic M -quasi-invertible Cantor system (Y, T) , with branch point $\mathbf{y}$ has an induced system (Y', S) where $\mathbf{y} \in Y'$. If (Y', S) is isomorphic to $(X_{\mathcal{B}'}, V_{\mathcal{B}'})$, then (Y, T) is isomorphic to $(X_{\mathcal{B}'}, V_{\mathcal{B}'})$ where $\mathcal{B}'$ is obtained from $\mathcal{B}$ by adding or removing a finite number of edges to $\mathcal{E}_1$, and changing the ordering on the affected vertices and edges.* ■

Let τ be a (not necessarily left proper) substitution with fixed point $\mathbf{u}$, and suppose that $u_0 = a$. A return word to a in $\mathbf{u}$ is a word $\mathbf{w}$ such that

1. u_0 is a prefix of $\mathbf{w}$;
2. There is no other occurrence of u_0 in $\mathbf{w}$;
3. $\mathbf{w}u_0 \in \mathcal{L}_{\tau}$.

This definition is a special case of the definition of a return word in [Dur98]. The set of return words $\mathcal{R}$ is finite, since $\mathbf{u}$ is almost periodic. This notion is the one-sided generalization of two sided return words in [DHS99], and we use their notation here. The set of return words $\mathcal{R}$ is finite, since $\mathbf{u}$ is almost periodic. This notion is the one-sided generalization of two sided return words in [DHS99], and we use their notation here. Ordering $\mathcal{R}$ according to the order of appearance of a return word in $\mathbf{u}$, we have a bijection ψ from $R := \{1, 2, \dots, |\mathcal{R}|\}$ to $\mathcal{R}$. The τ -fixed point $\mathbf{u}$, and so every element in X_{τ} , is a concatenation of return words, with possibly a suffix of a return word as a prefix. Extend ψ by concatenation to $\psi : R^{\mathbb{N}} \rightarrow [u_0] \subset \mathcal{A}^{\mathbb{N}}$. Since any $\mathbf{y} \in \psi(R^{\mathbb{N}})$ can only be partitioned in one way, using Property 2 of the definition of a return word, ψ is injective.

Let $\mathcal{D}(\mathbf{u}) \in R^{\mathbb{N}}$ be the unique sequence such that $\psi(\mathcal{D}(\mathbf{u})) = \mathbf{u}$. Let (Y, σ) be the subshift spanned by $\mathcal{D}(\mathbf{u}) \in R^{\mathbb{N}}$.

LEMMA 16. *(Y, σ) is isomorphic to the system induced by (X_{τ}, σ) on $[u_0]$ via the map ψ .*

Proof: For each $\mathbf{y} \in Y$, $\psi(\sigma \mathbf{y}) = \sigma^k(\psi(\mathbf{y}))$, where k is the length of the word in $\mathcal{R}$ corresponding to y_0 . So $\psi(\sigma^n(\mathbf{y})) \in [u_0]$ for all natural n , thus $\psi(Y) \subset [u_0]$. If $\mathbf{x} \in [u_0]$, then $\sigma^{n_k}(\mathbf{u}) \rightarrow \mathbf{x}$ for some $n_k \rightarrow \infty$. For large k , $\sigma^{n_k}(\mathbf{u}) \in [u_0]$, so $\sigma^{n_k}(\mathbf{u}) = \psi(\sigma^{j_k}(\mathcal{D}(\mathbf{u}))) \in \psi(Y)$ for some $j_k \rightarrow \infty$. Hence $\mathbf{x} \in \psi(Y)$. It follows that $\psi \circ \sigma = \sigma_{[u_0]} \circ \psi$ for $\mathbf{y} \in Y$. ■

The substitution τ can be used to define a substitution on R : if $\mathbf{w}$ is a return word, then $\mathbf{w}$ starts with u_0 , and since $\tau(u_0)$ starts with u_0 , so does $\tau(\mathbf{w})$. Also, $\mathbf{w}u_0$ is a word in $\mathbf{u}$, so $\tau(\mathbf{w}u_0) = \tau(\mathbf{w})u_0 \dots$ is also a word in $\mathbf{u}$. Hence $\tau(\mathbf{w})u_0 \dots$ is a concatenation of unique return words, and a prefix of some return word, so that $\tau(\mathbf{w})$ is a unique concatenation of return words. Thus if $j \in R$ corresponds to $\mathbf{w} \in \mathcal{R}$, and $\tau(\mathbf{w}) = \mathbf{w}_1 \mathbf{w}_2 \dots \mathbf{w}_r$, define $\tau_1(j) := i_1 i_2 \dots i_r$ where $i_k \in R$ corresponds to $\mathbf{w}_k \in \mathcal{R}$.

LEMMA 17. *Suppose τ have generating fixed point $\mathbf{u}$. Then*

1. τ_1 (or some power of τ_1) is left proper, primitive, aperiodic injective, with $\mathcal{D}(\mathbf{u})$ its fixed point;
2. τ_1 is recognizable; and
3. If τ is M -quasi-invertible, then τ_1 is M -quasi-invertible. Conversely, if τ is quasi-invertible, and τ_1 is M -quasi-invertible, then τ is M -quasi-invertible.

Proof:

1. Note that if $1 \in R$ corresponds to $\mathbf{w}$ in $\mathcal{R}$, then $\mathbf{w}u_0$ is a prefix of $\mathbf{u}$, and so is $\tau^n(u_0)$ for each n . Choose n large enough so that $|\tau^n(u_0)| > |\mathbf{w}|$, so that $\mathbf{w}u_0$ is a prefix of $\tau^n(u_0)$. If $j \in R$, and $\mathbf{w}' \in \mathcal{R}$ corresponds to j , then $\mathbf{w}'$ begins with u_0 , so that $\tau^n(u_0)$ and so $\mathbf{w}$ is a prefix of $\tau^n(\mathbf{w}')$. Hence 1 is the first letter of $\tau_1(j)$.

Since $\psi \circ \tau_1 = \tau \circ \psi$, then $\tau_1(D(\mathbf{u})) = D(\mathbf{u})$. Since τ is primitive, so is τ_1 . If $\tau_1(i) = \tau_1(j)$, then either $\tau(\alpha) = \tau(\beta)$ for some $\alpha \neq \beta$, contradicting injectivity of τ , or the conditions in Theorem 1 are satisfied, contradicting recognizability of τ .

2. Suppose that τ_1 is not recognizable. Then by Theorem 1, there exist letters i and j in R and some $\mathbf{y} \in Y$ with $\tau_1(i)$ a proper suffix of $\tau_1(j)$, and $\tau_1(i)\mathbf{y}$ and $\tau_1(j)\mathbf{y}$ appearing in Y with the same 1-cutting of $\mathbf{y}$. Suppose that $\psi(i) = \mathbf{w} = w_1 \dots w_l$ and $\psi(j) = \mathbf{w}' = w'_1 \dots w'_l$. Find the smallest k in $\{0 \dots l-1\}$ such that $w_{l-k} = w'_{l-k}$. Such a k exists since otherwise $\mathbf{w}$ is a suffix of $\mathbf{w}'$, and since $\mathbf{w} \neq \mathbf{w}'$, then $w_1 = u_0$ occurs in $w'_2 \dots w'_l$, a contradiction to Part 2 of the definition of a return word. Now as $\tau_1(i)$ is a proper suffix of $\tau_1(j)$, there exist words $\mathbf{v}$ and $\mathbf{v}'$ in R^+ , with $\mathbf{v}$ a suffix of $\mathbf{v}'$, $\mathbf{v}$ a suffix of $\tau_1(i)$, $\mathbf{v}'$ a suffix of $\tau_1(j)$, and words $\mathbf{p}, \mathbf{p}'$ and $\mathbf{s} \in \mathcal{A}^+$, with $\psi(\mathbf{v}) = \mathbf{p}\tau(w_{l-k})\mathbf{s}$ and $\psi(\mathbf{v}') = \mathbf{p}'\tau(w'_{l-k})\mathbf{s}$. If $|\tau(w_{l-k})| < |\tau(w'_{l-k})|$, then $\tau(w_{l-k})$ is a proper suffix of $\tau(w'_{l-k})$; and if $|\tau(w_{l-k})| > |\tau(w'_{l-k})|$, then $\tau(w'_{l-k})$ is a proper suffix of $\tau(w_{l-k})$. Finally if $|\tau(w_{l-k})| = |\tau(w'_{l-k})|$, then $\tau(w_{l-k}) = \tau(w'_{l-k})$, which is not possible as τ is injective on letters.

Thus (without loss of generality) $\tau(w_{l-k})$ is a proper subword of $\tau(w'_{l-k})$, and if $\mathbf{y}' = \tau(w_{l-k+1} \dots w_l)\psi(\mathbf{y})$, then $\tau(w_{l-k})\mathbf{y}'$ and $\tau(w'_{l-k})\mathbf{y}'$ appear in X_τ with the same 1-cutting, a contradiction to the recognizability of τ .

3. Note that if $\mathbf{x} \in X_\tau$ has distinct σ -preimages $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_M$, then in $[u_0]$, there exist $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_M$ with $\mathbf{y}_i = \sigma^{-t_i}\mathbf{x}_i$, where $-t_i$ is the first time a preimage of $\mathbf{x}_i$ is in $[u_0]$, - that this point is well-defined follows from the quasi-invertibility and the aperiodicity of (X_τ, σ) . Let $\mathbf{y} = \sigma_{[u_0]}(\mathbf{x})$ if $\mathbf{x} \notin [u_0]$, and $\mathbf{x} = \mathbf{y}$ otherwise. Then $\mathbf{y}$ has M distinct $\sigma_{[u_0]}$ -preimages, $\mathbf{y}_1, \dots, \mathbf{y}_M$. Since $\mathbf{x}$ is unique, so is $\mathbf{y}$. Now use the isomorphism in Lemma 16 to transfer this information to (Y, σ) .

Conversely, if (X_τ, σ) is N -quasi-invertible and $\mathbf{x}$ has N σ -preimages, $\{\mathbf{x}_1, \dots, \mathbf{x}_N\}$, then if t is the first re-entry time of $\mathbf{x}$ into $[u_0]$, $\sigma^t(\mathbf{x})$ has N $\sigma_{[u_0]}$ -preimages $\{\sigma^{-t_1}(\mathbf{x}_1), \dots, \sigma^{-t_N}(\mathbf{x}_N)\}$. Thus $\psi^{-1}(\sigma^t(\mathbf{x}))$ has N σ -preimages, and as (X_{τ_1}, σ) is M -quasi-invertible, $N = M$. ■

THEOREM 18. *Suppose that τ is M -quasi-invertible, with a generating fixed point $\mathbf{u}$, and branch point $\mathbf{y}$. Then there exists an ordered semi-proper, stationary Bratteli diagram $\mathcal{B}_{\tau^*}$ such that $(X_{\mathcal{B}}, V_{\tau^*})$ is topologically conjugate to (X_τ, σ) .*

Proof: Using the fixed point $\mathbf{u}$, and working with the induced system $([u_0], \sigma_{[u_0]})$, we use Lemma 17 to find τ_1 such that $([u_0], \sigma_{[u_0]})$ is conjugate to (X_{τ_1}, σ) . By Part 3 of Lemma 17, τ_1 is M -quasi-invertible. Now using Corollary 12, there is some τ^* left proper, and quasi-invertible, with its fixed point $\mathbf{u}^*$ as branch point, such that $(X_{\tau_1}, \sigma) = (X_{\tau^*}, \sigma)$ (If τ_1 's branch point is fixed, let $\tau_1 = \tau^*$). By Theorem 13 (X_{τ^*}, σ) , and so (X_{τ_1}, σ) , is isomorphic to $(X_{\mathcal{B}_{\tau^*}}, V_{\tau^*})$. If $\mathbf{y} \in [u_0]$, then by Lemma 14, (X_τ, σ) is a primitive of $([u_0], \sigma_{[u_0]})$, and $\mathbf{y}$ is mapped to the minimal element of $(X_{\mathcal{B}_{\tau^*}}, V_{\tau^*})$. Using Theorem 15, (X_τ, σ) is isomorphic to $(X_{\mathcal{B}}, V_{\tau^*})$ where $\mathcal{B}$ is obtained from $\mathcal{B}^*$ by the addition of a finite number of edges to E_1 , and V_{τ^*} is the corresponding adic map.

If $\mathbf{y} \notin [u_0]$, write (X_τ, σ) as a primitive over a sufficiently small cylinder set V containing $\mathbf{y}$, as in Lemma 14, and such that if a level of this partition $\mathcal{P}$ intersects $[u_0]$, then it is contained in $[u_0]$: thus $\mathcal{P}$ has $[u_0]$ split up as a union of cylinder sets defined by words of length N . We make V small enough so that no element in V returns to V before it passes through $[u_0]$. Telescope just the first M levels of the Bratteli diagram $\mathcal{B}_{\tau^*}$, so that the elements in $\mathcal{P}$ which are subsets of $[u_0]$ are represented by edges from v_0 to the first level. Now add levels in E_1 according to how the cylinder sets in $[u_0]$ appear in the clopen partition

of X_τ . For example, If C and C' are cylinder sets contained in $[u_0]$ and appearing as $C = V_k^i$ and $C' = V_k^{i'}$ with $i < i'$ and no set $V_k^{i+1}, \dots, V_k^{i'-1}$ intersects $[u_0]$, then in the telescoped Bratteli diagram, the edge e' corresponding to C' is the successor of the edge e corresponding to C ; now insert $i' - i - 1$ new edges between e and e' . The fact that elements in V have to pass through $[u_0]$ before returning to V means that you can add enough levels to the telescoped stationary Bratteli diagram (which is still defined by τ^*) so that the new Bratteli diagram $\mathcal{B}_{\tau^*}$ generates an adic system $(X_{\mathcal{B}^*}, V_{\mathcal{B}^*})$ isomorphic to the original (X_τ, σ) . ■

Lemma 9 in [For97] is used in the two sided version of this previous result, to show that (aperiodic, primitive) substitution subshifts are conjugate to a stationary adic system where there are only single edges from the vertex v_0 to any vertex in $\mathcal{V}_1$. Although the statement of this lemma is also true for semi-proper Bratteli diagrams, it is not clear that the resulting ordered stationary diagram corresponds to a recognizable, or injective substitution. See for example, Figure 4 in [DHS99].

Examples:

9. If τ is suffix permutative, with one fixed point $\mathbf{u}$, then it is quasi-invertible, and (X_τ, σ) is a primitive over (X_{τ_1}, σ) , as $\phi(\mathbf{u}) \in X_{\tau_1}$.
10. If $\tau(a) = aac$, $\tau(b) = bcc$, and $\tau(c) = abc$, then (X_τ, σ) is 3-quasi-invertible, with the non-invertible element $\mathbf{y}$ satisfying $c\tau(\mathbf{y}) = \mathbf{y}$. Since all τ -words have a proper common suffix, Corollary 12 tells us that $(X_\tau, \sigma) = (X_{\tau^*}, \sigma)$ where $\tau^*(\mathbf{y}) = \mathbf{y}$. Now apply Theorem 13.
11. If $\tau(a) = aac$, $\tau(b) = bcc$, $\tau(c) = adbc$ and $\tau(d) = adbd$, then since $da \notin \mathcal{L}_\tau$ and $cb \notin \mathcal{L}_\tau$, both fixed points are σ -invertible. Inspection of maximal proper common suffixes of the τ -substitution words of letters in any subset $\mathcal{A}^*$ of $\mathcal{A}$ leads to the existence of only one branch point $\mathbf{y}$ satisfying $c\tau(\mathbf{y}) = \mathbf{y}$. Lemma 10 cannot directly apply because the family of τ -substitution words does not have a common suffix. If we consider instead the induced system $([c], \sigma_{[c]})$, (X_τ, σ) is a primitive over $([c], \sigma_{[c]})$. Define $\tau^*(a) = caa$, $\tau^*(b) = c$, $\tau^*(c) = cadb$ and $\tau^*(d) = cadbdb$. One can now prove, similarly to the proof of Lemma 11 to show that $\tau^*(\mathbf{y}) = \mathbf{y}$. Thus $([c], \sigma_{[c]})$ is itself a (left proper, quasi-invertible) substitution subshift, and so has a stationary adic representation $(X_{\mathcal{B}^*}, V_{\mathcal{B}^*})$. Now Theorem 15 applies, so that (X_τ, σ) also has a stationary adic representation. In this example it is relatively straightforward to find the right τ^* ; making the adic representation of (X_τ, σ) more straightforward than if we had followed the proof of Theorem 18; in general though it is not clear how a branch point $\mathbf{y}$ can be seen as the fixed point of some τ^* which is obtained directly from the definition of τ .
12. *Minimal rank one subshifts* are defined by substitutions on $\mathcal{A} = \{0, 1\}$ of the form

$$\tau(0) = 0^{n_1} 1^{m_1} 0^{n_2} 1^{m_2} \dots 0^{n_{k-1}} 1^{m_{k-1}} 0^{n_k} \text{ and } \tau(1) = 1,$$

where $k < \infty$, and n_1 and n_k are positive. The latter condition ensures that the resulting substitution subshift is minimal, with generating fixed point $\mathbf{u} := \lim_{n \rightarrow \infty} \tau^n(0)$. These systems are equivalent to rank-one systems defined by ‘cutting and stacking’ where there are a bounded number of cuts and spacers added, and the same cutting-and-stacking rule is obeyed at each stage. A comprehensive exposition of rank one systems is given in [Fer97]. Suppose $n_i \neq n_j$ for some i, j , so that τ is aperiodic. Then (X_τ, σ) is quasi-invertible if and only if $m_1 = m_2 = \dots = m_{k-1} = m > 0$. For, the generating fixed point $\mathbf{u}$ is a branch point, and the fixed point $1^{m_i} \mathbf{u}$ is another branch point if and only if $m_i < m_j$ for some j .

If $\tau(0) = 0^{n_1}1^m0^{n_2}1^m \dots 0^{n_{k-1}}1^m0^{n_k}$, then there are two return words to 0, $w_1 = 0$ and $w_2 = 01^m$. In this case τ_1 is defined on $\{1, 2\}$ and $\tau_i = wi$ where $w = 1^{n_1-1}21^{n_2-1}2 \dots 1^{n_k-1}$. By Lemma 17, τ_1 is left proper and 2-quasi-invertible. Thus (X_τ, σ) is a primitive of $([0], \sigma_{[0]})$ and by Theorem 18, τ has a semi proper adic representation by adding $m-1$ edges from v_0 to the vertex in $\mathcal{V}_1$ corresponding to the letter '1' in the Bratteli representation of τ_1 . For $1 \leq k \leq M$, the τ -fixed points $1^k u$ live on (different) levels of the tower over (X_τ, σ) .

If for some i, j , $m_i \neq m_j$, then τ_1 is still of the form $\tau_1(i) = wi$, where $w = 1^{n_1-1}21^{n_2-1}3 \dots 1^{n_{k-1}-1}k1^{n_k-1}$, so that τ_1 is M-quasi-invertible, with $M \geq 3$. Thus (X_{τ_1}, σ) is quasi-invertible, even though (X_τ, σ) is not, and we cannot generate an adic representation for τ from one for τ_1 . For example, if $\tau(0) = 00100110$, then the return words are $\mathcal{R} = \{0, 01, 011\}$ and τ_1 is 3-almost periodic, with x^1, x^2 , and x^3 the preimages of $\mathcal{D}(u)$. In X_τ though, $\phi(x^1)$, and $\sigma(\phi(x^2))$ are preimages of u , and $\phi(x^2)$ and $\sigma(\phi(x^3))$ are preimages of $\sigma(\phi(x^2))$. So an adic representation of (X_τ, σ) would need 4 maximal elements, one of which is also minimal, and another minimal element.

13. For any primitive substitution τ whose composition matrix has a rational Perron-Frobenius eigenvalue, there exists a quasi-invertible primitive substitution τ' whose associated two sided substitution system $(X_{\tau'}^{\mathbb{Z}}, \sigma)$ is orbit equivalent to $(X_\tau^{\mathbb{Z}}, \sigma)$. To see this we use the results in Corollary 6.7 and Theorem 6.15 in [Yua02]. First, given τ , with Perron value λ , we find a positive integer d so that $\langle \mu(E) : E \subset X_\tau \text{ is clopen} \rangle = \{ \frac{n}{m} : n \in \mathbb{Z}, m \text{ is a factor of } d \cdot \lambda^n \text{ for some } n \}$, where the former is order isomorphic to the dimension group modulo the infinitesimal subgroup of (X_τ, σ) . Then given d and λ , both greater than 1, the author describes conditions on the composition matrix of τ' so that $\langle \mu(E) : E \subset X_{\tau'}, E \text{ clopen} \rangle = \{ \frac{n}{m} : n \in \mathbb{Z}, m \text{ is a factor of } d \cdot \lambda^n \text{ for some } n \}$. In particular if the substitution τ' defined on d letters has the substitution matrix

$$\begin{pmatrix} \lambda^m & \lambda^m & \dots & \lambda^m & (\lambda^m - d + 1)\lambda^m \\ \lambda^m & \lambda^m & \dots & (\lambda^m - d + 1)\lambda^m & \lambda^m \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ (\lambda^m - d + 1)\lambda^m & \lambda^m & \dots & \lambda^m & \lambda^m \end{pmatrix}$$

then τ' has the desired dimension group (modulo the infinitesimal subgroup) We claim here that τ' can be chosen to be quasi invertible. In particular if τ is defined by $\tau'(1) = 1^{\lambda^m}2^{\lambda^m}3^{\lambda^m} \dots (d-1)^{\lambda^m}d^{(\lambda^m-d)\lambda^m}d^{\lambda^m}$, $\tau'(2) = 1^{\lambda^m}2^{\lambda^m}3^{\lambda^m} \dots (d-1)^{\lambda^m}(d-1)^{(\lambda^m-d)\lambda^m}d^{\lambda^m}$, $\dots \tau'(d) = 1^{\lambda^m}2^{\lambda^m}3^{\lambda^m} \dots (d-1)^{\lambda^m}(1)^{(\lambda^m-d)\lambda^m}d^{\lambda^m}$, then τ' is quasi invertible.

14. Here is an example of a quasi-invertible substitution whose two-sided subshift conjugacy class is not the same as its one-sided subshift conjugacy class. Define $\tau(a) = aabaa$, $\tau(b) = abcab$, and $\tau(c) = aabac$. The fixed point $u = a \dots$ is a 3-branch point, and $(X_\tau^{\mathbb{N}}, \sigma)$ is quasi-invertible. Let $u_L = \dots u_{-2}u_{-1}$ be the left infinite sequence generated by $\{\tau^n(a)\}$, v_L the one generated by $\{\tau^n(b)\}$, and w_L the one generated by $\{\tau^n(c)\}$. Note that if $n \neq -1$, $u_n = w_n$. The words of length 3 in $\mathcal{L}_\tau$ are $\mathcal{W}_3 = \{aab, aba, baa, aaa, abc, bca, cab, bac, aca, caa\}$. Define a local rule $\phi : \mathcal{W}_3 \rightarrow \{\alpha, \beta, \gamma, d, e, f, g, h\}$ with left and right radius one, as $\phi(aaa) = \phi(baa) = \phi(aca) = \alpha$, $\phi(aba) = \beta$, $\phi(caa) = \gamma$, and let ϕ map the remaining words in $\mathcal{W}_3$ in a one-one fashion to $\{d, e, f, g, h\}$. Let Φ be the shift-commuting factor mapping corresponding to ϕ and Let $Y = \Phi(X_\tau^{\mathbb{Z}})$.

LEMMA 19. *The map $\Phi : X_\tau^{\mathbb{Z}} \rightarrow Y$ is injective.*

Proof: Assume that for some $\mathbf{x} \neq \mathbf{y}$, $\Phi(\mathbf{x}) = \Phi(\mathbf{y})$. So there is some n such that $x_{n-1}x_nx_{n+1} \neq y_{n-1}y_ny_{n+1}$, and yet both words belong to $\{baa, aaa, aca\}$. There are three cases to consider, all similar to prove, so we look at the case when $x_{n-1}x_nx_{n+1} = baa$ and $y_{n-1}y_ny_{n+1} = aaa$. If baa appears in a sequence here, it either appears as a subword of $\tau(a) = aabaa$, or $\tau(ba) = abcabaabaa$ or $\tau(bc) = abcabaabac$. On the other hand aaa can only appear as a subword of $\tau(aa) = aabaaaabaa$ or $\tau(ac) = aabaaaabac$ or $\tau(ab) = aabaaabcab$. This means that $x_{n-2}x_{n-1}x_n = aba$ and $y_{n-2}y_{n-1}y_n = aaa$ or baa . Either way $\Phi(\mathbf{x})_n = \beta$ and $\Phi(\mathbf{y})_n = \alpha$, contradicting $\Phi(\mathbf{x}) = \Phi(\mathbf{y})$. $\blacksquare$

Now Theorem 4 in [DHS99] tells us that (Y, σ) is either a substitution system, or a stationary odometer. Since (X_τ, σ) is not an odometer, and Φ is a conjugacy, so (Y, σ) is generated by some substitution τ' . Hence the 2-sided subshifts generated by τ and τ' are conjugate.

If $\mathbf{x} = x_0x_1 \dots$ is a one sided sequence, let $\Phi(x) = \phi(x_0x_1x_2)\phi(x_1x_2x_3) \dots$. Note that τ' has two 2-branch points. For,

$$\begin{aligned} \Phi(\mathbf{u}_L \cdot \mathbf{u}) &= \Phi(\dots u_{-2}u_{-1})\phi(u_{-2}u_{-1}u_0) \cdot \phi(u_{-1}u_0u_1)\Phi(\mathbf{u}) = \\ \Phi(\dots u_{-2}u_{-1})\phi(aaa) \cdot \phi(aaa)\Phi(\mathbf{u}) &= \Phi(\dots u_{-2}u_{-1})\alpha \cdot \alpha\Phi(\mathbf{u}) \end{aligned}$$

and

$$\begin{aligned} \Phi(\mathbf{w}_L \cdot \mathbf{u}) &= \Phi(\dots w_{-2}w_{-1})\phi(w_{-2}w_{-1}u_0) \cdot \phi(w_{-1}u_0u_1)\Phi(\mathbf{u}) = \\ \Phi(\dots w_{-2}w_{-1})\phi(aca) \cdot \phi(caa)\Phi(\mathbf{u}) &= \Phi(\dots w_{-2}w_{-1})\alpha \cdot \gamma\Phi(\mathbf{u}), \end{aligned}$$

so that $\Phi(\mathbf{u})$ is a 2-branch point; while

$$\begin{aligned} \Phi(\mathbf{v}_L \cdot \mathbf{u}) &= \Phi(\dots v_{-2}v_{-1})\phi(v_{-2}v_{-1}u_0) \cdot \phi(v_{-1}u_0u_1)\Phi(\mathbf{u}) = \\ \Phi(\dots v_{-2}v_{-1})\phi(aba) \cdot \phi(baa)\Phi(\mathbf{u}) &= \Phi(\dots v_{-2}v_{-1})\beta \cdot \alpha\Phi(\mathbf{u}), \end{aligned}$$

so that $\alpha\Phi(\mathbf{u})$ is also a 2-branch point. Thus the one sided substitution systems generated by τ and τ' cannot be topologically conjugate.

6. (N, n) -substitutions.

In this section we consider substitutions which have more than one branch point. The next theorem is a special case of Theorem 21, which we include as its proof is simpler:

THEOREM 20. *Suppose that τ is defined on $\mathcal{A} = \{0, 1, \dots, n-1\}$, and is prefix and suffix permutative, so that there are n fixed points, the i -th of which is a k_i -branch point. If $\mathcal{L}_\tau$ has N words of length 2, then there exists τ' defined on $\mathcal{A}' = \{0, 1, \dots, N-1\}$ such that τ' is suffix permutative, with n fixed points, where each is a k_n branch point and where $(X_\tau, \sigma) \cong (X_{\tau'}, \sigma)$, which is topologically conjugate to $(X_{\mathcal{B}_{\tau'}}, V_{\tau'})$.*

Proof: First note that all branch points are τ -fixed. By taking a power of τ if necessary we can assume that $\tau(i)$ begins and ends with i . Ordering the words of length two in $\mathcal{L}_\tau$ lexicographically, define the right radius one local rule ϕ on these words by mapping the j -th element in this list to j , and let Φ be the shift commuting map associated with ϕ . Note that the image of ϕ is $\mathcal{A}' := \{0, \dots, N-1\}$ where N is the number of words of length 2 in $\mathcal{L}_\tau$ and satisfies $N = \sum_{i=0}^{n-1} k_i$. Now define the substitution τ' on $\mathcal{A}'$ as follows: if $a = \phi(ij)$, then $\tau'(a) := \phi(\tau(i))a$. It is clear that τ' is suffix permutative, with n fixed points. We will show that (X_τ, σ) is topologically conjugate to $(X_{\tau'}, \sigma)$ by showing that $\mathbf{v}_i := \Phi(\tau^\infty(i))$ is the τ' -fixed in $(X_{\tau'}, \sigma)$. If that is the case, then since Φ is injective and shift commuting, and the fixed points generate the subshift, we are done.

To see this we show by induction that $\tau^n(a) = \phi(\tau^n(i))a$ for all n , whenever $\phi(ij) = a$. For $n = 1$ the assertion is true by definition. Assuming the case true for n , we have

$$\begin{aligned}
\tau^{n+1}(a) &= \tau'^n(\tau'(a)) = \tau'^n(\phi(\tau(i))a) = \tau'^n(\phi(ii_1 \dots i_{l_i} i)a) \\
&= \tau'^n(\phi(ii_1)\phi(i_1 i_2) \dots \phi(i_{l_i} i)a) \\
&= \tau'^n(\phi(ii_1))\tau'^n(\phi(i_1 i_2)) \dots \tau'^n(\phi(i_{l_i} i))\tau'^n(a) \\
&= \phi(\tau^n(i))\phi(ii_1)\phi(\tau^n(i_1))\phi(i_1 i_2) \dots \phi(\tau^n(i_{l_i}))\phi(i_{l_i} i)\phi(\tau^n(i))a \\
&\stackrel{*}{=} \phi(\tau^n(i)\tau^n(i_1) \dots \tau^n(i_{l_i})\tau^n(i))a \\
&= \phi(\tau^{n+1}(i))a,
\end{aligned}$$

where $*$ follows since τ is prefix and suffix permutative. Now apply this to $\mathbf{v}_i \in \phi[ni + i_1]$.

To see that τ' has an adic representation, note that if $i' = \phi(ij)$ generates a τ' -fixed point, then the k_i predecessors of i' are $\{\phi(\ell i)\}$ as ℓ ranges over all predecessors of i in $\mathcal{L}_\tau$. As for distinct i , these sets are disjoint, Proposition 9 implies that $V_{\tau'}$ is continuous, and Proposition 10 that $(X_{\tau'}, \sigma) \cong (X_{\mathcal{B}_{\tau'}}, V_{\tau'})$. ■

Examples:

1. If $\tau(0) = 01$ and $\tau(1) = 10$, then τ' is defined as $\tau'(0) = 1320$, $\tau'(1) = 1321$, $\tau'(2) = 2012$, and $\tau'(3) = 2013$. Note that $\mathcal{B}_{\tau'}$ has four maximal elements $M_0, \dots, M_3$ and two minimal elements m_1, m_2 , where M_i is constant in $\mathcal{B}_\tau$ through vertex i . The words of length 2 in $\mathcal{L}_{\tau'}$ are $\{01, 12, 13, 20, 21, 32\}$ so $V_{\mathcal{B}} : X_{\mathcal{B}} \rightarrow X_{\mathcal{B}}$ can be defined continuously with $V_{\mathcal{B}}(M_0) = V_{\mathcal{B}}(M_2) = m_1$ and $V_{\mathcal{B}}(M_1) = V_{\mathcal{B}}(M_3) = m_2$.
2. Let $\tau(0) = 0120$, $\tau(1) = 100101121$, $\tau(2) = 20212$. Then $\tau^\infty(0)$ and $\tau^\infty(1)$ are 3-branch points, and $\tau^\infty(2)$ is a 2-branch point. Here $N = 8$, so define τ' on $\{0, 1, 2, \dots, 7\}$ by $\tau'(0) = 1560$, $\tau'(1) = 1561$, $\tau'(2) = 1562$, $\tau'(3) = 301314573$, $\tau'(4) = 301314574$, $\tau'(5) = 3013141575$, $\tau'(6) = 62756$, and $\tau'(7) = 62757$. One can check that $\tau'^\infty(1)$ is a 3-branch point with predecessors $0\tau'^\infty(1)$, $3\tau'^\infty(1)$, and $6\tau'^\infty(1)$; that $\tau'^\infty(3)$ is also a 3-branch point with predecessors $1\tau'^\infty(3)$, $4\tau'^\infty(3)$ and $7\tau'^\infty(3)$, and that $\tau'^\infty(6)$ is a 2-branch point with predecessors $2\tau'^\infty(6)$ and $7\tau'^\infty(6)$. Thus extend $V_{\mathcal{B}_{\tau'}}$ by mapping $V_{\mathcal{B}_{\tau'}}(\mathbf{M}_0) = V_{\mathcal{B}_{\tau'}}(\mathbf{M}_3) = V_{\mathcal{B}_{\tau'}}(\mathbf{M}_6) = \mathbf{m}_0$, $V_{\mathcal{B}_{\tau'}}(\mathbf{M}_1) = V_{\mathcal{B}_{\tau'}}(\mathbf{M}_4) = V_{\mathcal{B}_{\tau'}}(\mathbf{M}_7) = \mathbf{m}_1$ and $V_{\mathcal{B}_{\tau'}}(\mathbf{M}_2) = V_{\mathcal{B}_{\tau'}}(\mathbf{M}_5) = \mathbf{m}_2$.

THEOREM 21. *Suppose that τ is defined on $\mathcal{A} = \{0, 1, \dots, n-1\}$, so that there are m right-fixed points $\{\mathbf{p}_i : 1 \leq i \leq m\}$, each of which is a k_i -branch point. Suppose there are no other branch points. If $\mathcal{L}_\tau$ has N words of length 2, then there exists τ' defined on the N -letter alphabet $\mathcal{A}'$ such that*

1. $(X_\tau, \sigma) \cong (X_{\tau'}, \sigma)$,
2. τ' has m right-fixed points $\{\mathbf{m}_i = (m_i, \dots) : 1 \leq i \leq m\}$,
3. τ' has $\sum_{i=1}^m k_i$ left-fixed points,
4. $V_{\tau'}$ is continuous, and
5. $(X_{\tau'}, \sigma) \cong (X_{\mathcal{B}_{\tau'}}, V_{\tau'})$

Proof: Assume that the fixed point $\mathbf{p}_i$ starts with the letters $p_i p_i^1$. We can assume, by taking a power of τ if necessary, that all substitutions words start with $p_i p_i^1$ for some $i \in \{1, \dots, m\}$, and end with a letter which generates a left infinite τ -fixed point. We define a right-sided, radius one factor mapping $\phi : \mathcal{L}_2(\tau) \rightarrow \mathcal{A}'$ where $\mathcal{A}'$ is an N -letter alphabet. Let $\phi(p_i, p_i^1) := m_i$. If $\mathbf{sp}_i$ is a σ -predecessor of $\mathbf{p}_i$, where the cylinder set $[sp_i]$ has not appeared as the home of one of the right infinite τ -fixed points, define $\phi(sp_i) = M_l$ for some l , as p_i

and s range over all allowed letters. For the remaining words ab in $\mathcal{L}_2(\tau)$, let ϕ map them injectively to letters that are distinct from the already specified m_i 's and M_i 's. Let $\mathcal{A}'$ be the resulting alphabet. Define τ' on $\mathcal{A}'$ as follows. If $i \in \mathcal{A}$, let p^i be the first letter of $\tau(i)$, and s^i the last letter of $\tau(i)$. If $a = \phi(ij)$, let $\tau'(a) := \phi(\tau(i))\phi(s^i p^j)$. Here $\phi(\tau(i))$ is a word of length $|\tau(i)| - 1$. The following two claims are straightforward to verify, starting with the definition of τ' .

Claim 1: If $\alpha\beta$ is a 2-word appearing in $\tau'(\gamma)$, where $\alpha = \phi(ij)$ and $\beta = \phi(i'j')$, then the suffix of $\tau'(\alpha)$ is $s = \phi(s^i p^{i'})$, as is the suffix of $\tau'^n(\alpha)$ for each $n \in \mathbb{N}$.

Claim 2: If $\alpha = \phi(i_\alpha, j_\alpha)$ for $\alpha \in \mathcal{A}'$, and $\tau'(\alpha) = \alpha_1 \alpha_2 \dots \alpha_k$, then $\tau(i_\alpha) = i_{\alpha_1} i_{\alpha_2} \dots i_{\alpha_k}$.

Claim 3: If $\alpha = \phi(i_\alpha, j_\alpha)$ is a letter in $\mathcal{A}'$, then $(\tau')^n(\alpha) \setminus \text{a suffix} = \phi(\tau^n(i))$ for each $n \in \mathbb{N}$. If $n = 1$ then it is true by definition.

If it is true for n , then

$$\begin{aligned}
 (\tau')^{n+1}(\alpha) &= (\tau')^n(\tau'(\alpha)) = (\tau')^n(\alpha_1 \alpha_2 \dots \alpha_k) \\
 &= (\tau')^n(\alpha_1) (\tau')^n(\alpha_2) \dots (\tau')^n(\alpha_k) \\
 &\stackrel{\text{Claim 1}}{=} \phi(\tau^n(i_{\alpha_1})) \phi(s^{i_{\alpha_1}} p^{i_{\alpha_2}}) \phi(\tau^n(i_{\alpha_2})) \phi(s^{i_{\alpha_2}} p^{i_{\alpha_3}}) \dots \phi(\tau^n(i_{\alpha_k})) \ell \\
 &= \phi(\tau^n(i_{\alpha_1} i_{\alpha_2} \dots i_{\alpha_k})) \ell \stackrel{\text{Claim 2}}{=} \phi(\tau^n(\tau(i_\alpha))) \ell = \phi(\tau^{n+1}(i_\alpha)) \ell,
 \end{aligned}$$

where ℓ is some letter in $\mathcal{A}'$. The result follows.

Thus in particular Φ maps the τ -fixed points $\{\mathbf{p}_i\}$ to the τ' -fixed points $\{\mathbf{m}_i\}$ and since ϕ is 1-1, Φ is shift commuting, and the fixed points generate the subshift, then (X_τ, σ) is topologically conjugate to $(X_{\tau'}^\mathbb{N}, \sigma)$. Statements 2 and 3 follow by definition of τ' , and to see 4, note that if $m_k \neq m_l$, then $i_{m_k} \neq i_{m_l}$. Thus if $Mm_k \in \mathcal{L}_{\tau'}$, then $Mm_i \notin \mathcal{L}$ for any other m_i . Thus M can be followed by at most one of the m'_i s, and must be followed by at least one, by minimality of τ .

Clearly τ' is primitive. We show that τ' is recognizable and $\tau' : X_{\tau'} \rightarrow X_{\tau'}$ is a homeomorphism (here τ' is not injective, so that Part 4 of Proposition 2 cannot be directly applied). If $\mathbf{x}', \mathbf{y}' \in X_{\tau'}$, let the Φ -preimages of $\mathbf{x}'$ and $\mathbf{y}'$ in X_τ be $\mathbf{x}$ and $\mathbf{y}$ respectively. If $\tau'(\mathbf{x}') = \tau'(\mathbf{y}')$, then using Part 3 of Proposition 2, we have $|\tau'(x'_n)| = |\tau'(y'_n)|$ for each $n \geq 0$, so that $\tau'(x'_n) = \tau'(y'_n)$ for each $n \geq 0$. Now $\tau'(x'_n) = \phi(\tau(x_n))\phi(s_{x_n} p_{x_{n+1}})$ and $\tau'(y'_n) = \phi(\tau(y_n))\phi(s_{y_n} p_{y_{n+1}})$, and since τ is injective, if $x_n \neq y_n$, either $\phi(\tau(x_n)) \neq \phi(\tau(y_n))$, or $s_{x_n} \neq s_{y_n}$. Thus $\mathbf{x} = \mathbf{y}$ and so $\mathbf{x}' = \mathbf{y}'$. If τ' is not recognizable, then there exist letters a', b' and for each L , words w'_L satisfying the conditions of Theorem 1. This implies that there exist corresponding letters a, b such that $\tau(b)$ is a proper suffix of $\tau(a)$, and, as in the proof of injectivity of τ' the words w'_L lift to words $w_L \in \mathcal{L}_\tau$ such that conditions of Theorem 1 are satisfied for τ contradicting recognizability of τ .

Thus all branch points of τ' are fixed points, and using Corollary 8, the partitions $(\mathcal{P}'_n)$ generate the topology of $X_{\tau'}$. Now Proposition 10 implies that $(X_{\tau'}, \sigma) \cong (X_{\mathcal{B}_{\tau'}}, V_{\tau'})$. ■

Example: If $\tau(0) = 01200$, $\tau(1) = 11201$ and $\tau(2) = 10121$, then τ has two fixed points, both of which are 2-branch points. Define $\phi(01) = m_1$, $\phi(11) = m_2$, $\phi(00) = M_1$, $\phi(10) = M_2$, $\phi(12) = a$, $\phi(20) = b$ and $\phi(21) = c$. If $A_i := \phi(\tau(i))$, then $A_0 = m_1 a b M_1$, $A_1 = m_2 a b m_1$ and $A_2 = M_2 m_1 a c$; and τ' is defined as $\tau'(M_1) = A_0 M_1$, $\tau'(m_1) = A_0 m_1$, $\tau'(M_2) = A_1 M_2$, $\tau(m_2) = \tau(a) = A_1 m_2$, $\tau(b) = A_2 M_2$ and $\tau(c) = A_2 m_2$. Note τ' has four left infinite fixed points, generated by $\{m_1, m_2, M_1, M_2\}$, and the predecessors of m_1 are $\{M_1, M_2\}$, while those of m_2 are $\{m_1, m_2\}$.

REFERENCES

- [AR91] Pierre Arnoux and Gérard Rauzy. Représentation géométrique de suites de complexité $2n + 1$. *Bull. Soc. Math. France*, 119(2):199–215, 1991.
- [BDH03] Marcy Barge, Beverly Diamond, and Charles Holton. Asymptotic orbits of primitive substitutions. *Theoret. Comput. Sci.*, 301(1-3):439–450, 2003.
- [BKM09] S. Bezuglyi, J. Kwiatkowski, and K. Medynets. Aperiodic substitution systems and their Bratteli diagrams. *Ergodic Theory Dynam. Systems*, 29(1):37–72, 2009.
- [BKMY11] Sergey Bezuglyi, Jan Kwiatkowski, Konstantin Medynets, and Reem Yassawi. Orderings on bratteli diagrams and vershik maps. *preprint*, 2011.
- [CK71] Ethan M. Coven and Michael S. Keane. The structure of substitution minimal sets. *Trans. Amer. Math. Soc.*, 162:89–102, 1971.
- [CMPS93] D. Crisp, W. Moran, A. Pollington, and P. Shiue. Substitution invariant cutting sequences. *J. Théor. Nombres Bordeaux*, 5(1):123–137, 1993.
- [DHS99] F. Durand, B. Host, and C. Skau. Substitutional dynamical systems, Bratteli diagrams and dimension groups. *Ergodic Theory Dynam. Systems*, 19(4):953–993, 1999.
- [Dur98] F. Durand. A characterization of substitutive sequences using return words. *Discrete Mathematics*, 179:89–101, 1998.
- [Fer97] Sébastien Ferenczi. Systems of finite rank. *Colloq. Math.*, 73(1):35–65, 1997.
- [For97] A. H. Forrest. K -groups associated with substitution minimal systems. *Israel J. Math.*, 98:101–139, 1997.
- [GPS95] Thierry Giordano, Ian F. Putnam, and Christian F. Skau. Topological orbit equivalence and C^* -crossed products. *J. Reine Angew. Math.*, 469:51–111, 1995.
- [Hos86] B. Host. Valeurs propres des systèmes dynamiques définis par des substitutions de longueur variable. *Ergodic Theory Dynam. Systems*, 6(4):529–540, 1986.
- [HPS92] Richard H. Herman, Ian F. Putnam, and Christian F. Skau. Ordered Bratteli diagrams, dimension groups and topological dynamics. *Internat. J. Math.*, 3(6):827–864, 1992.
- [HZ01] Charles Holton and Luca Zamboni. Directed graphs and substitutions. *preprint*, 2001.
- [Mos92] Brigitte Mossé. Puissances de mots et reconnaissabilité des points fixes d’une substitution. *Theoret. Comput. Sci.*, 99(2):327–334, 1992.
- [PS73] Karl Petersen and Leonard Shapiro. Induced flows. *Trans. Amer. Math. Soc.*, 177:375–390, 1973.
- [Que87] Martine Queffélec. *Substitution dynamical systems—spectral analysis*, volume 1294 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1987.
- [Yua02] H. Yuasa. On the topological orbit equivalence in a class of substitution minimal systems. *Tokyo J. Math.*, 25(2):221–240, 2002.